

ON δ -HOMOGENEOUS RIEMANNIAN MANIFOLDS

V.N. BERESTOVSKIĬ AND YU.G. NIKONOROV

ABSTRACT. We study in this paper previously defined by V.N. Berestovskii and C.P. Plaut δ -homogeneous spaces in the case of Riemannian manifolds. Every such manifold has non-negative sectional curvature. The universal covering of any δ -homogeneous Riemannian manifold is itself δ -homogeneous. In turn, every simply connected Riemannian δ -homogeneous manifold is a direct metric product of an Euclidean space and compact simply connected indecomposable homogeneous manifolds; all factors in this product are itself δ -homogeneous. We find different characterizations of δ -homogeneous Riemannian spaces, which imply that any such space is geodesic orbit (g.o.) and every normal homogeneous Riemannian manifold is δ -homogeneous. The g.o. property and the δ -homogeneity property are inherited by closed totally geodesic submanifolds. Then we find all possible candidates for compact simply connected indecomposable Riemannian δ -homogeneous non-normal manifolds of positive Euler characteristic and a priori inequalities for parameters of the corresponding family of Riemannian δ -homogeneous metrics on them (necessarily two-parametric). We prove that there are only two families of possible candidates: non-normal (generalized) flag manifolds $SO(2l+1)/U(l)$ and $Sp(l)/U(1) \cdot Sp(l-1)$, $l \geq 2$, investigated earlier by W. Ziller, H. Tamaru, D.V. Alekseevsky and A. Arvanitoyeorgos. At the end we prove that the corresponding two-parametric family of Riemannian metrics on $SO(5)/U(2) = Sp(2)/U(1) \cdot Sp(1)$ satisfying the above mentioned (strict!) inequalities, really generates δ -homogeneous spaces, which are not normal and are not naturally reductive with respect to any isometry group.

2000 Mathematical Subject Classification: 53C20 (primary), 53C25, 53C35 (secondary).

Key words and phrases: homogeneous spaces, homogeneous spaces of positive Euler characteristic, geodesic orbit spaces, Clifford-Wolf translations, geodesics, normal homogeneous Riemannian spaces, submetry, Riemannian submersion.

CONTENTS

1. Introduction	2
2. Preliminaries	4
3. General properties of δ -homogeneous spaces	5
4. Totally geodesic submanifolds	11
5. Additional symmetries of δ -homogeneous metrics	14
6. δ -vectors	16
7. On the topology of compact homogeneous spaces	17
8. Homogeneous spaces of positive Euler characteristic	19
9. On algebraic corollaries of the δ -homogeneity	23
10. On δ -homogeneous manifold of one special type	24
11. Root systems of compact simple Lie algebras	26
12. On the group G_2	28
13. Calculations with roots	28
14. The special second case	32
15. On the space $SO(5)/U(2) = Sp(2)/U(1) \cdot Sp(1) = \mathbb{C}P^3$	35
References	39

The first author is supported in part by RFBR (grant N 04-01-00315-a). The second author is supported in part by RFBR (grant N 05-01-00611-a) and by the Council on grants of the President of Russian Federation for supporting of young russian scientists and leading scientific schools of Russian Federation (grants NSH-8526.2006.1 and MD-5179.2006.1).

1. INTRODUCTION

Historically, the assembly of homogeneous Riemannian manifolds under considerations has been gradually extended and about thirty years ago it included all such manifolds. Nevertheless, the division of them into particular classes is very important.

We shall mention here classes, which can be characterized by some properties of their isometry groups with connection to their geodesics. B. Riemann separated all manifolds of constant sectional curvature, which are characterized by the property of free movability of figures. Later E. Cartan introduced and classified all *symmetric* Riemannian manifolds. Then K. Nomizu introduced and studied *naturally reductive* homogeneous Riemannian manifolds which include as special cases symmetric spaces and *normal homogeneous* manifolds. The latter have non-negative sectional curvatures and include all symmetric spaces with nonnegative sectional curvatures. A little later A. Selberg introduced another generalization of symmetric spaces, namely, *weakly symmetric* Riemannian spaces. At last, *geodesic orbit (g.o.)* homogeneous Riemannian manifolds have been discovered. This class includes properly as special all previously mentioned classes, see [31], [52]. Every simply connected Riemannian g.o. manifold of dimension ≤ 5 is naturally reductive [31]. Riemannian g.o. manifolds of dimension 6, which are not naturally reductive, are classified in [31]; for recent results in dimension 7 we refer to [20]. A. Selberg proved that every weakly symmetric Riemannian manifold M is *commutative*, i.e. it admits a transitive motion Lie group G such that the Lie algebra of G -invariant differential operators on M is commutative [39]. If G is connected and $M = G/H$, the latter is equivalent to the property that the functional space $L^1(H \backslash G/H)$ is commutative, i.e. (G, H) is a *Gelfand pair*, or the property that for every unitary irreducible representation of G , the dimension of H -fixed set is ≤ 1 , i.e. (G, H) is a *spherical pair*. J. Lauret obtained an example of commutative non weakly symmetric Riemannian manifold [33]. On the other hand, if (G, H) is a spherical pair with compact simple Lie group G and its closed subgroup H , then G/H is weakly symmetric [36]. A classification of such pairs is known from [32], [36], [5]. Let us remark that besides symmetric spaces, there is no complete classification of manifolds in other classes, mentioned above, although normal homogeneous manifolds do not require in some sense such classification.

We prove in this paper that the previously defined in [8] δ -homogeneous spaces constitute in the case of Riemannian manifolds a new class of homogeneous manifolds situated between normal homogeneous and g.o. manifolds. These manifolds, unlike all previously mentioned classes, have very simple, purely metric definition, which really can be applied to any metric space. Namely, an arbitrary metric space (M, ρ) is called *δ -homogeneous*, if for every two points $x, y \in M$ there is an isometry f of (M, ρ) onto itself, which moves x to y and has the maximal displacement at the point x , i.e. $f(x) = y$ and $\rho(x, f(x)) \geq \rho(z, f(z))$ for all $z \in M$. If we can always take such a motion f from an isometry group G of (M, ρ) , then (M, ρ) is called *G - δ -homogeneous*. In the Riemannian case we shall take as G only connected transitive Lie groups.

The consideration and methods in this paper go from general to more and more specific.

In Section 2 we bring main definitions, earlier results, and simple examples. In particular, any Lie group with a bi-invariant Riemannian metric or any direct metric product of δ -homogeneous spaces is δ -homogeneous. Every δ -homogeneous Riemannian manifold has non-negative sectional curvature (Proposition 1).

In Section 3 we get general results on δ -homogeneous Riemannian manifolds (M, μ) . If (M, μ) is G - δ -homogeneous and G normalizes a closed subgroup H of the full isometry group $I(M)$ of (M, μ) , then the quotient (orbit) space $H \backslash M$ with the quotient Riemannian metric is δ -homogeneous (Theorem 3). As a corollary (1), we get that every normal homogeneous Riemannian manifold is δ -homogeneous. Then we prove that the universal locally isometric covering of (M, μ) is δ -homogeneous (Corollary 2); (M, μ) is either compact or it is isometric to a direct metric product of an Euclidean space and some compact δ -homogeneous Riemannian manifold (Theorem 4). Since any homogeneous Riemannian manifold (M, μ) is

an orbit space of its universal covering $(\tilde{M}, \tilde{\mu})$ by central discrete subgroup Γ in the (unit) connected component of $I(\tilde{M})$, where Γ is isomorphic to $\pi_1(M)$, then previous results imply that the study of δ -homogeneous Riemannian manifolds entirely reduces to the simply connected compact case. Then we get four useful necessary and sufficient conditions for a (homogeneous) connected Riemannian manifolds (M, μ) to be δ -homogeneous; we will mention here two of them. First: $(M = G/H, \mu)$ with the corresponding inner metric ρ is G - δ -homogeneous if and only if it is G -normal in the generalized sense (Corollary 4). The latter means that there is a bi-invariant Finsler (inner) metric F on G such that the natural projection $p : (G, F) \rightarrow (G/H, \rho)$ is submetry (see Definition 3). Notice that in the case when F is Riemannian (inner), (M, μ) would be G -normal. Second: $(M = G/H, \mu)$ is G - δ -homogeneous if and only if every geodesic γ in (M, μ) is an orbit of a 1-parameter motion group of (M, μ) in G , generated by a Killing vector field, attaining a maximal value of its length at γ (Theorem 7). As a corollary (6), every δ -homogeneous Riemannian manifold is geodesic orbit (g.o.). At the same time, g.o. Lobachevski's space of constant negative curvature cannot be δ -homogeneous by Proposition 1.

In Section 4 we prove that every closed totally geodesic submanifold of a δ -homogeneous (respectively, g.o.) Riemannian manifold is δ -homogeneous (respectively, g.o.) itself, see Theorem 11 (respectively, 12). As a corollary (Theorem 13), every factor of a δ -homogeneous or g.o. direct metric product has the same property. By all previous results, the study of all δ -homogeneous Riemannian spaces reduces to the compact simply connected indecomposable case; we can separate further the cases of zero or positive Euler characteristic. In the second half of the section we find some algebraic properties of geodesic vectors on a homogeneous Riemannian space $(G/H, \mu)$, i.e. vectors in the Lie algebra $\mathfrak{g}$ of the Lie group G , that tangent to a 1-parameter subgroup in G with geodesic orbit through the point $H \in (G/H, \mu)$. Besides later applications, we use them to prove that if $(G/H, \mu)$ is G - δ -homogeneous and L is a Lie subgroup of G such that $H \subset L \subset G$, then L/H with the metric, induced by μ , is δ -homogeneous.

In Section 5 we find additional isometries of δ -homogeneous Riemannian manifolds and find some applications of them.

Similarly to geodesic vectors, we define a δ -vector on $(G/H, \mu)$ as a vector in $\mathfrak{g}$ that tangent to (the unique) right-invariant vector field Y on G with the property that the Killing vector field $X = dp(Y)$ on $(G/H, \mu)$ has maximal value of its length at the point $H \in (G/H, \mu)$. Remark that every δ -vector is a geodesic vector. In Section 6 we find general properties of δ -vectors. We use essentially these properties later. The space $(G/H, \mu)$ is G - δ -homogeneous if and only if every vector $v \in T_H(G/H)$ can be represented in the form $v = dp(w)$ for some δ -vector w .

In Section 7 we give (mainly known) results on compact simply connected homogeneous spaces $M = G/H$, in particular, Hopf-Samelson Theorem 17, which implies that $\chi(M) \geq 0$ and characterizes the case, when $\chi(M) > 0$, by the condition $\text{rk}(G) = \text{rk}(H)$. Let us mark Theorem 20, which states that every proper Lie subalgebra $\mathfrak{h}$ of the Lie algebra $\mathfrak{g}$ of a simple compact connected Lie group G , containing the Lie algebra $\mathfrak{t}$ of a maximal torus $T \subset G$, is the Lie algebra of the unique closed connected Lie subgroup $H \subset G$. Moreover, $M = G/H$ is a simply connected compact connected homogeneous space of positive Euler characteristic. This gives an algebraic description of all simply connected compact indecomposable homogeneous Riemannian manifolds (M, μ) with $\chi(M) > 0$ by the Kostant's Theorem 22.

In Section 8 we give known results on compact simply connected homogeneous spaces of positive Euler characteristic. Also we prove Theorem 25 which implies that every naturally reductive homogeneous Riemannian manifolds of positive Euler characteristic is normal (hence, δ -homogeneous).

In Section 9 algebraic corollaries of δ -homogeneity of the first and the second order are found in Theorem 26. Note that the first order condition is simply the condition for geodesic vectors.

In the next sections we consider only compact simply connected homogeneous spaces of positive Euler characteristic.

In Section 10 we find some algebraic identities and inequalities for δ -homogeneous manifolds of one special type. As we shall prove in Section 13, any (compact simply connected) indecomposable δ -homogeneous non-normal Riemannian manifold (M, μ) with $\chi(M) > 0$ must have such type. Especially important are Propositions 21, 22, and 27.

In Section 11 is given necessary information on roots and structural constants of compact simple Lie algebras with respect to their Killing forms and Cartan subalgebras. Mark especially the identity (11.17).

We show in Section 12 that all G_2 - δ -homogeneous Riemannian metrics on homogeneous spaces with positive Euler characteristic are normal.

Observe that roots of (simple compact) Lie algebras A_l , D_l , e_6 , e_7 and e_8 have one and the same length, while the roots of Lie algebra g_2 , B_l , C_l and f_4 have two different lengths. One knows also that the Weyl group of any simple Lie algebra acts transitively on the set of roots with equal lengths. With the help of these facts and the identity (11.17) we prove in Section 13 that the set of G - δ -homogeneous Riemannian metrics on G/H with $\chi(G/H) > 0$ and compact simple Lie group G is one- or two-parametric; we have necessarily the first case (that is, only G -normal metrics), if the Lie algebra $\mathfrak{g}$ has roots of equal length, as for Lie groups $G = SU(l+1)$, $SO(2l)$, E_6 , E_7 , E_8 (Proposition 32 and Corollary 13). We shall have only G -normal metrics also in the case, when $H = T$ (Proposition 14). In the case of two-parametric family we get with the help of Proposition 27 a priori inequalities (13.21) for these parameters in Proposition 33.

Further investigations of two-parametric case in Section 14 shows that possible candidates one can find only among flag manifolds $SO(2l+1)/U(l)$ and $Sp(l)/U(1) \cdot Sp(l-1)$, where $l \geq 2$. All invariant metrics on these manifolds are weakly symmetric (hence, g.o.) [52]. Moreover, among (generalized) flag manifolds only $SO(2l+1)/U(l)$ and $Sp(l)/U(1) \cdot Sp(l-1)$, $l \geq 2$, admit non-normal invariant g.o. metrics [4].

Using Proposition 21 and spectra of matrices, we prove in Section 15 that two-parametric family of Riemannian metrics on $SO(5)/U(2)$, which satisfies the inequalities (13.21), really give us $SO(5)$ - δ -homogeneous spaces. The limiting cases of this inequalities represent $SO(5)$ -normal and $SO(6)$ -normal spaces respectively, while all other metrics are $SO(5)$ - δ -homogeneous and non-normal (Theorem 30). We are planning to investigate all other possible cases, mentioned in the previous paragraph, separately.

Some unsolved questions are posed in different places of the text.

The first author is very obliged to Mathematics Department of University of Tennessee, Knoxville, USA, for hospitality and visiting position while a part of this paper have been prepared.

2. PRELIMINARIES

Definition 1. Let (X, d) be a metric space and $x \in X$. An isometry $f : X \rightarrow X$ is called a $\delta(x)$ -translation (a Clifford-Wolf translation), if x is a point of maximal displacement of f , i.e. for every $y \in X$ the relation $d(y, f(y)) \leq d(x, f(x))$ holds (respectively, f displaces all points of (X, d) the same distance, i.e. $d(y, f(y)) = d(x, f(x))$ for every $y \in X$).

Definition 2. A metric space (X, d) is called (G) - δ -homogeneous (respectively, (G) -Clifford-Wolf homogeneous), if for every $x, y \in X$ there exists a $\delta(x)$ -translation (respectively, Clifford-Wolf translation) of (X, d) (from an isometry group G), moving x to y .

It is clear that any Clifford-Wolf translation is a $\delta(x)$ -translation for every point $x \in X$, any (G) -Clifford-Wolf homogeneous space is (G) - δ -homogeneous, and the latter one is (G) -homogeneous.

Example 1. Every Lie group with a bi-invariant inner metric (G, r) and every odd-dimensional Euclidean sphere (of the unit radius) $S^{2n+1} \subset \mathbb{E}^{2(n+1)}$ with the induced inner (Riemannian) metric is Clifford-Wolf homogeneous space. In the first case it is enough to use left

translations on some fixed element of the group. The second statement is proved essentially by Clifford himself, which explains the term in Definition 2.

Example 2. One can easily see that a direct metric product of δ -(respectively, Clifford-Wolf) homogeneous spaces is again δ -(respectively, Clifford-Wolf) homogeneous.

In the paper [8] the following results are obtained.

Theorem 1 (Berestovskii-Plaut [8]). *Every locally compact δ -homogeneous space of curvature bounded below in the sense of Alexandrov has non-negative curvature.*

Theorem 2 (Berestovskii-Plaut [8]). *Every non-compact locally compact homogeneous inner metric space of non-negative curvature in the sense of Alexandrov is isometric to a direct metric product of finite dimensional Euclidean space and a compact homogeneous inner metric space of non-negative curvature.*

Remark 1. In the Riemannian case, the last theorem easily follows from Toponogov's theorem in [44], which states that every complete Riemannian manifold (M, μ) with nonnegative sectional curvature, containing a metric line, is isometric to a direct Riemannian product $(N, \nu) \times \mathbb{R}$. Later J. Cheeger and D. Gromoll in [17] generalized Toponogov's theorem to complete Riemannian manifolds of nonnegative Ricci curvature.

If (M, μ) is a Riemannian manifold with inner metric ρ , then Theorem 1 implies

Proposition 1. *Every δ -homogeneous Riemannian manifold (M, ρ) has non-negative sectional curvature.*

Definition 3. A map of metric spaces $f : (M, r) \rightarrow (N, q)$ is called a *submetry*, if it maps every closed ball $B(x, s) \subset (M, r)$ with the radius s and the center x onto the closed ball $B(f(x), s) \subset (N, q)$ with the radius s and the center $f(x)$, [9].

Note that a smooth map of complete Riemannian spaces is submetry if and only if it is a Riemannian submersion [9].

Definition 4. A locally compact inner metric (respectively, Riemannian) space $(M = G/H, \rho)$ with a transitive locally compact topological (respectively, Lie) group G and a stabilizer subgroup H at a point $x \in M$ is called *G -normal in generalized (respectively, usual) sense*, if G admits a bi-invariant (respectively, Riemannian bi-invariant) inner metric r such that the natural projection $(G, r) \rightarrow (G/H, \rho)$ is a submetry.

3. GENERAL PROPERTIES OF δ -HOMOGENEOUS SPACES

Definition 5. An inner metric space (M, ρ) is called *restrictively (G) - δ -homogeneous* (respectively, *restrictively (G) -Clifford-Wolf homogeneous*) if for every $x \in M$ there exists a number $r(x) > 0$ such that for every two points y, z in the open ball $U(x, r(x))$ there exists a $\delta(y)$ -translation (respectively, a Clifford-Wolf translation) of the space (M, ρ) (from the isometry group G), moving y to z . The supremum $R(x)$ of all such numbers $r(x)$ is called *the (G) - δ -homogeneity radius* (respectively, *the (G) -Clifford-Wolf homogeneity radius*) of the space (M, ρ) at the point x .

Proposition 2. *Every restrictively (G) - δ -homogeneous locally compact complete inner metric space is (G) - δ -homogeneous.*

Proof. It is clear that (in the notation of Definition 5) the function $R(x)$, $x \in M$, is equal identically to $+\infty$, i.e. the space (M, ρ) is (G) - δ -homogeneous, or it satisfies the inequality $|R(x_1) - R(x_2)| \leq \rho(x_1, x_2)$. In the last case the function $R(x)$, $x \in M$, is positive and continuous.

Let us consider arbitrary points x, y of a metric space (M, ρ) , and suppose that this space satisfies the above-stated condition. Then one can join the points x and y by some shortest $[x, y]$. According to the above discussion, one can divide sequentially this shortest

by points $x_0 = x, x_1, \dots, x_m = y$ such that for every l , where $0 \leq l \leq m-1$, there exists a $\delta(x_l)$ -translation f_l of the space (M, ρ) (from the group G), moving the point x_l to the point x_{l+1} . Now the triangle inequality implies that the composition $f := f_{m-1} \circ \dots \circ f_0$ is a $\delta(x)$ -translation of the space (M, ρ) (from the group G), moving the point x to the point y . ■

Theorem 3. *Let (M, r) be a locally compact inner metric space which is G - δ -homogeneous. Suppose that the group G normalizes some closed subgroup H of the full isometry group $\text{Isom}(M)$ of M (supplied by the compact-open topology). Then the quotient (orbit) space $H \backslash M$ with the quotient metric ρ is a (G) - δ -homogeneous (locally compact inner metric) space.*

Proof. According to S.E. Cohn-Vossen theorem [18], every complete locally compact inner metric space is finitely-compact, i.e., every its closed bounded subset is compact. It is proved in the paper [6] that any closed subgroup of the full isometry group (with the compact-open topology) of arbitrary finitely-compact space has closed orbits. This implies that the group H has closed orbits in M .

On the ground of this fact it is easy to prove that the canonical projection $p : (M, r) \rightarrow (H \backslash M, \rho)$ is a submetry. This is equivalent to the following two properties:

- 1) the map p does not increase distances;
- 2) for every three points $x, y \in H \backslash M$, $\xi \in p^{-1}(x)$, there exists a point $\eta \in p^{-1}(y)$ such that $r(\xi, \eta) = \rho(x, y)$.

Now let us consider arbitrary points $x, y \in H \backslash M$ and the corresponding points ξ, η from Property 2). By condition there is a $\delta(\xi)$ -translation F of the space (M, r) from the group G such that $F(\xi) = \eta$. Since the group G normalizes the group H , there is an isometry f of the space $(H \backslash M, \rho)$, induced by the isometry F . Moreover, $f(x) = p(F(\xi)) = p(\eta) = y$. Now for any point $z = p(\zeta) \in H \backslash M$ Properties 1) and 2) imply the relations

$$\begin{aligned} \rho(x, f(x)) &= \rho(x, y) = r(\xi, \eta) = r(\xi, F(\xi)) \geq \\ r(\zeta, F(\zeta)) &\geq \rho(p(\zeta), p(F(\zeta))) = \rho(z, f(z)), \end{aligned}$$

i.e. f is a $\delta(x)$ -translation of the space $(H \backslash M, \rho)$ moving the point x to the point y . Therefore, the space $(H \backslash M, \rho)$ is G - δ -homogeneous. ■

Corollary 1. *Every (G) -normal in the generalized sense homogeneous locally compact inner metric space is (G) - δ -homogeneous. As a corollary, any (G) -normal (maybe, in the generalized sense) homogeneous Riemannian manifold is (G) - δ -homogeneous.*

Proof. Let a (G) -normal (in the generalized sense) homogeneous space under consideration be a (metric) quotient space $(G/H, \rho)$ of a locally compact topological group (G, r) with a bi-invariant inner metric r by its compact subgroup H . Then the group of left translations of the group (G, r) is a transitive group of Clifford-Wolf translations, and it commutes with the group of right translations by elements of the subgroup H which consists of some isometries of the space (G, r) . Now it is enough to use Theorem 3. ■

Proposition 3. *The universal locally isometric covering of a δ -homogeneous (respectively, a restrictively Clifford-Wolf homogeneous) Busemann's G -space is a δ -homogeneous (respectively, a restrictively Clifford-Wolf homogeneous) Busemann's G -space.*

Proof. Busemann's G -spaces are defined in his book [16].

Let $p : (\tilde{M}, \tilde{\rho}) \rightarrow (M, \rho)$ be the universal locally isometric covering map for a δ -homogeneous (respectively, a restrictively Clifford-Wolf homogeneous) Busemann's G -space (M, ρ) . It is clear that $(\tilde{M}, \tilde{\rho})$ is a Busemann's G -space. By Theorem 28.10 in [16], the group G of all motions of the space $(\tilde{M}, \tilde{\rho})$, which cover motions of the space (M, ρ) , is transitive on $\tilde{M}$, and the group Γ of deck transformations of the covering p is a normal subgroup of the group G . Therefore, there is a number $r > 0$ such that the map p is isometry on every open ball $U(x, r) \subset (\tilde{M}, \tilde{\rho})$.

According to Proposition 2, it is enough to show that the space $(\tilde{M}, \tilde{\rho})$ is restrictively δ -homogeneous (respectively, restrictively Clifford-Wolf homogeneous). Consider arbitrary points x, y in $(\tilde{M}, \tilde{\rho})$ with the condition $\tilde{\rho}(x, y) < r$. Since (M, ρ) is δ -homogeneous (respectively, restrictively Clifford-Wolf homogeneous), there is a $\delta(p(x))$ -translation (respectively, a Clifford-Wolf translation) f of the space (M, ρ) such that $f(p(x)) = p(y)$. From the above discussion we get that there is the unique map F of the space $(\tilde{M}, \tilde{\rho})$ onto itself covering the map f such that $F(x) = y$. It is clear that F is an isometry of the space $(\tilde{M}, \tilde{\rho})$ and also a $\delta(x)$ -translation (respectively, a Clifford-Wolf translation). This means that the space $(\tilde{M}, \tilde{\rho})$ is restrictively δ -homogeneous (respectively, restrictively Clifford-Wolf homogeneous). ■

Corollary 2. *The universal Riemannian covering of a δ -homogeneous (respectively, a restrictively Clifford-Wolf homogeneous) Riemannian manifold is δ -homogeneous (respectively, restrictively Clifford-Wolf homogeneous).*

Lemma 1. *Suppose that the Riemannian manifold (M, μ) is isometric to the direct metric product $(K, \mu_1) \times (\mathbb{E}^m, \mu_2)$, where (K, μ_1) is a compact homogeneous Riemannian manifold, and $(\mathbb{E}^m, \mu_2)$ is a finite dimensional Euclidean space. Then every isometry f of the space (M, μ) has the form $f = f_1 \times f_2$, where f_1 (respectively, f_2) is an isometry of the space (K, μ_1) (respectively, $(\mathbb{E}^m, \mu_2)$).*

Proof. It is easy to see that a geodesic in (M, μ) is a metric line if and only if it is situated in some Euclidean subspace $\{k\} \times \mathbb{E}^m$. Therefore, any isometry f of the space (M, μ) transposes such subspaces. Since f keeps the orthogonality, f must transpose also all fibers of the form $K \times \{e\}$. This proves Lemma. ■

Lemma 2. *If $M = M_1 \times M_2$ is a direct product of Riemannian manifolds, then every its isometry of the form $f = f_1 \times f_2$ is a $\delta(x)$ -translation for the point $x = (x_1, x_2) \in M$ if and only if both isometries $f_1 : M_1 \rightarrow M_1$ and $f_2 : M_2 \rightarrow M_2$ are δ -translations at the points $x_1 \in M_1$ and $x_2 \in M_2$ respectively.*

Proof. Let us remind that

$$\rho((x_1, x_2), (y_1, y_2)) = \sqrt{\rho_1^2(x_1, y_1) + \rho_2^2(x_2, y_2)},$$

where ρ, ρ_1, ρ_2 are inner metrics of spaces M, M_1, M_2 respectively. This easily implies the sufficiency. Suppose that $f = f_1 \times f_2$ is a $\delta(x)$ -translation of the space M at the point $x = (x_1, x_2)$, but, for instance, f_1 is not a δ -translation at the point x_1 . Then there is a point x'_1 such that $\rho_1(x'_1, f_1(x'_1)) > \rho_1(x_1, f_1(x_1))$. Therefore,

$$\begin{aligned} \rho((x_1, x_2), f(x_1, x_2)) &= \sqrt{\rho_1^2(x_1, f_1(x_1)) + \rho_2^2(x_2, f_2(x_2))} < \\ &\sqrt{\rho_1^2(x'_1, f_1(x'_1)) + \rho_2^2(x_2, f_2(x_2))} = \rho((x'_1, x_2), f(x'_1, x_2)), \end{aligned}$$

which contradicts to assumptions of Lemma. ■

Theorem 4. *Any δ -homogeneous Riemannian manifold (M, g) is either compact, or it is isometric to the direct metric product of an Euclidean space and some compact δ -homogeneous Riemannian manifold.*

Proof. This theorem immediately follows from Proposition 1, Theorem 2, Lemma 1 and Lemma 2. ■

From Theorem 4 and Proposition 3 we immediately obtain

Corollary 3. *The universal Riemannian covering $(\tilde{M}, \tilde{\mu})$ of a δ -homogeneous compact Riemannian manifold (M, μ) is compact if and only if $\pi_1(M)$ is finite. In the opposite case $(\tilde{M}, \tilde{\mu})$ is isometric to a nontrivial direct metric product of a compact simply connected δ -homogeneous Riemannian space and an Euclidean space.*

Theorem 5. *A homogeneous space $M = G/H$ of a connected Lie group G by its compact subgroup H admits an invariant Riemannian δ -homogeneous metric if and only if G/H admits an invariant Riemannian metric of non-negative sectional curvature.*

Proof. The necessity follows from Proposition 1.

Let us prove the sufficiency. Suppose that $M = G/H$ admits an invariant Riemannian metric μ of non-negative sectional curvature.

If M is compact, then the Lie group G is compact and it admits a bi-invariant Riemannian metric γ . Then there is an unique Riemannian metric ν on M such that the canonical projecture $p : (G, \gamma) \rightarrow (M, \nu)$ is a Riemannian submersion. Moreover, ν is invariant on G/H , and $(G/H, \nu)$ is a G -normal homogeneous Riemannian manifold. According to Corollary 1, $(G/H, \nu)$ is a δ -homogeneous space.

Suppose, that M is noncompact. Then by Theorem 2, all assumptions of Lemma 1 are fulfilled, moreover, (K, μ_1) has non-negative sectional curvature. Therefore, the Lemma 1 is valid. Obviously, the set of all isometries of the type $\{f_1 | f = (f_1, f_2) \in G\}$ forms a precompact transitive isometry group G_1 of the compact space (K, μ_1) (relatively to the compact-open topology) with the closure $\Gamma_1 := \overline{G_1}$, which is a compact effective transitive isometry Lie group of the space (K, μ_1) . Consequently, the manifold K admits a Γ_1 -invariant Riemannian metric γ_1 such that (K, γ_1) is a normal homogeneous space of the Lie group Γ_1 . According to Corollary 1, (K, γ_1) is a δ -homogeneous space. The last reasonings imply that the Riemannian metric $g_0 = \gamma_1 \times \mu_2$ on M is invariant under the action of the group G . In this case the Riemannian manifold $(M, \mu_0) = (K, \gamma_1) \times (\mathbb{E}^m, \mu_2)$ is a δ -homogeneous space as a direct metric product of δ -homogeneous spaces. ■

Theorem 6. *Let (M, μ) be a smooth connected compact Riemannian manifold with inner metric ρ , and G be the identity component of the full isometry group of (M, μ) . Then the function $d : G \times G \rightarrow \mathbb{R}$ defined by the formula*

$$d(g, h) = \max_{x \in M} \rho(g(x), h(x)), \quad (3.1)$$

determines a bi-invariant metric on G compatible with its compact-open topology. In this case (G, d) is locally isometric to (G, D) for some bi-invariant inner metric D on G . Under identification of the Lie algebra G_e of the group G with the Lie algebra of Killing vector fields on (M, μ) , D coincides with the bi-invariant Finsler metric on G , determined by the $\text{Ad}(G)$ -invariant norm $\|\cdot\|$ on G_e , defined by the formula

$$\|X\| = \max_{x \in M} \sqrt{\mu(X(x), X(x))}. \quad (3.2)$$

Proof. One can check directly the bi-invariance of the metric d . The compactness of (M, μ) implies the compactness of the Lie group G . Then, since G is connected, the exponential map of the Lie algebra G_e to G is surjective.

Let $g \neq e$ be arbitrary element in G . Then $g = \exp(X)$ for some suitable Killing vector field X on (M, μ) . Let

$$\|X\| = \max_{x \in M} \sqrt{\mu(X(x), X(x))} = \sqrt{\mu(X(y), X(y))}.$$

According to Proposition 5.7 of Chapter VI in [28], the curve $\gamma(t) = \exp(tX)(y)$, $0 \leq t \leq 1$, is a segment of a geodesic in (M, μ) with the length $\|X\|$. It is known that for any other point $x \in M$ the curve $\exp(tX)(x)$, $0 \leq t \leq 1$, is parameterized proportionally to the arc-length with the coefficient of proportionality $\sqrt{\mu(X(x), X(x))}$, which does not exceed $\|X\|$. Therefore, the length of any arc of the second curve does not exceed the length of the corresponding arc of the geodesic γ .

The injectivity radius of the compact smooth manifold (M, μ) is bounded below by some number $r > 0$. If $0 \leq s\|X\| \leq r$; $t, s \in [0, 1]$, then it implies that for $g(s) = \exp(sX)$, $g(t) = \exp(tX)$, the point $\gamma(t)$ is the point of maximal displacement on (M, ρ) for the motion $g(s)$, since $\rho(g(s)(\gamma(t)), \gamma(t)) = s\|X\|$ according to equalities

$$g(s)(\gamma(t)) = g(s)(g(t)(y)) = g(s+t)(y) = \gamma(s+t).$$

Hence, $d(g(t), g(t+s)) = s\|X\|$, the length of the curve $g(t)$, $0 \leq t \leq 1$, in (G, d) equals to $\|X\|$. Therefore, one can join any two point in (G, d) by a curve of finite length (with respect to the metric d). Let D be the inner metric corresponding to d .

There exists a positive number s_0 such that $\exp : g \rightarrow G$ is a homeomorphism of some open subset V of g , containing the zero, onto the open ball $U(e, s_0)$ with the radius s_0 in (G, d) . Then the above reasonings imply that the curve $g(t)$, $0 \leq t \leq 1$, is a geodesic in (G, D) , and $D(g, h) = d(g, h)$, if $d(g, h) < \min(r, s)$. Also, $d \leq D$.

From the above calculations of the length of the geodesic $g(t) = \exp(tX)$, $0 \leq t \leq 1$, in (G, D) , it is clear that D is the bi-invariant Finsler (inner) metric on G determined by the $\text{Ad}(G)$ -invariant norm $\|\cdot\|$ on G_e , which defined by the formula (3.2). It is easy to check that this formula defines some norm on G_e . ■

Question 1. *Whether the metrics d and D coincide on G ?*

Theorem 7. *Let (M, μ) be a compact homogeneous Riemannian manifold. Then there exists a positive number $s > 0$ such that for arbitrary motion f of the space (M, g) with maximal displacement δ , which is less than s , there is unique Killing vector field X on (M, g) such that $\max_{x \in M} \sqrt{\mu(X(x), X(x))} = 1$ and $\gamma_X(\delta) = f$, where $\gamma_X(t)$, $t \in \mathbb{R}$ is the one-parameter motion group in (M, g) generated by the field X . If also f is a Clifford-Wolf translation, then the Killing field X has constant unit length on (M, μ) .*

Proof. Let us supply the identity component G of the full isometry group of (M, g) with the bi-invariant metric d as in Theorem 6. There is sufficiently small number $s > 0$ (which we can suppose smaller than the injectivity radius r of the manifold (M, μ)) such that the exponential map $\exp : g \rightarrow G$ is a homeomorphism of some neighborhood V of the zero in g onto an open ball $U(e, s)$ in (G, d) . Then for every motion f of the space (M, μ) with the condition $d(f, e) = \delta < s$ there exists the unique vector $Y \in V$ such that $\exp(Y) = f$. It was shown in the proof of Theorem 6 that for all such motions f we have $D(f, e) = d(f, e)$. This common value is equal also to the length of the path $\exp(\tau Y)$, $0 \leq \tau \leq 1$, which joins elements e and f , with respect to the bi-invariant norm $\|\cdot\|$ on TG from Theorem 6, and to the length $\|Y\|$. By the definition, $\|Y\| = \max_{x \in M} \sqrt{\mu(Y(x), Y(x))}$. Now it is clear that $X = (1/\delta)Y$ is an desired vector. The uniqueness of X follows from the above arguments.

Let us suppose also that f is a Clifford-Wolf translation. By the above construction we have

$$\|X\| = 1 = \max_{x \in M} \sqrt{\mu(X(x), X(x))} = \sqrt{\mu(X(x_1), X(x_1))} \quad (3.3)$$

for some point $x_1 \in M$. We state that

$$\sqrt{\mu(X(x), X(x))} \equiv 1.$$

Indeed, in the opposite case there would be a point $x_0 \in M$ such that $\sqrt{\mu(X(x_0), X(x_0))} = \varepsilon < 1$. Then the path $c(t) = \exp(tX)(x_0)$, $0 \leq t \leq \delta$, joins the point x_0 with the point $f(x_0)$ and has the length $\delta\varepsilon$. Therefore,

$$\rho(x_0, f(x_0)) \leq \delta\varepsilon < \delta = \rho(x_1, f(x_1)),$$

because, according to the condition (3.3), the orbit of the point x_1 under the action of the one-parameter group $\exp(tX)$, $t \in \mathbb{R}$, is a geodesic [28], and $\delta < r$. But this contradicts to the fact that f is a Clifford-Wolf translation. ■

Theorem 8. *Let (M, μ) be a compact connected (G) - δ -homogeneous Riemannian manifold with inner metric ρ , and let G be a closed connected (Lie) subgroup of the full isometry group of (M, μ) , supplied by the bi-invariant inner metric D as in Theorem 6 (more exactly, by its restriction to G). Then D is an inner bi-invariant metric on G . Let us fix a point $x_0 \in M$ and define a projection $p : G \rightarrow M$ by the formula $p(g) = g(x_0)$ such that under usual identification of M with G/H , where H is the stabilizer of G at the point x_0 , p coincides with the canonical projection $p : G \rightarrow G/H$. Then the map $p : (G, D) \rightarrow (M, \rho)$ is a submetry.*

Proof. The first statement easily follows from arguments in the last two paragraphs in the proof of Theorem 6, applied to G .

Now it is enough to check the properties 1) and 2) from the proof of Theorem 3.

1) Let $g, h \in G$. Then

$$\rho(p(g), p(h)) = \rho(g(x_0), h(x_0)) \leq \max_{x \in M} \rho(g(x), h(x)) = d(g, h) \leq D(g, h),$$

i.e. p does not increase distances.

2) Consider any points x, y in M and put $\rho(x, y) = a$. Let us choose arbitrary shortest K in (M, ρ) joining points x and y ; consider a geodesic $\gamma(s)$, $s \in \mathbb{R}$, in (M, μ) parameterized by the arc-length such that $\gamma(0) = x$, $\gamma(a) = y$ and $\gamma(s) \in K$, $0 \leq s \leq a$. Since (M, ρ) is G - δ -homogeneous, there is $\delta(x)$ -translation $g_t \in G$ of (M, ρ) , moving the point x to the point $\gamma(t)$, $0 < t \leq a$. Now if t is small enough, then by Theorems 6 and 7, there is an one-parameter group of motions $g(s) = \gamma_X(s) \in G$, $s \in \mathbb{R}$, such that $g(t) = g_t$ and $\max_{y \in M} \sqrt{\mu(X(y), X(y))} = \sqrt{\mu(X(x), X(x))}$. Then $g(s)(x) = \gamma(s)$, $s \in \mathbb{R}$.

Therefore, $D(e = g(0), g(s)) = d(e, g(s)) = s$ for $0 \leq s \leq a$. Suppose that $p(h) = h(x_0) = x$ for some element $h \in G$. Then

$$y = \gamma(a) = g(a)(x) = g(a)(h(x_0)) = p(g(a)h)$$

and

$$D(h, g(a)h) = D(e, g(a)) = a = \rho(x, y).$$

■

On the ground of Corollary 1 and Theorem 8 we obtain

Corollary 4. *A compact connected Riemannian manifold is (G) - δ -homogeneous if and only if it is (G) -normal in the generalized sense.*

Let us consider a compact Riemannian homogeneous manifold $(G/H, \mu)$, some $\text{Ad}(G)$ -invariant inner product $\langle \cdot, \cdot \rangle$ on the Lie algebra $\mathfrak{g}$ of the group G , the corresponding $\langle \cdot, \cdot \rangle$ -orthogonal direct sum decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ ($\mathfrak{h}$ is the Lie algebra of H), and $\text{Ad}(H)$ -invariant inner product $(\cdot, \cdot)$ on $\mathfrak{p}$ which defines the Riemannian metric μ . Then we can state the previous corollary as follows:

Theorem 9. *A compact Riemannian manifold $(G/H, \mu)$ is G - δ -homogeneous for Lie group G if and only if there exists an $\text{Ad}(G)$ -invariant centrally symmetric (relative to zero) convex body B in $\mathfrak{g}$ such that*

$$P(B) = \{v \in \mathfrak{p} \mid (v, v) \leq 1\},$$

where $P : \mathfrak{g} \rightarrow \mathfrak{p}$ is $\langle \cdot, \cdot \rangle$ -orthogonal projection. One can take $C = \{w \in \mathfrak{g} \mid \|w\| \leq 1\}$ as B .

Corollary 5. *The vector space $\mathfrak{p}$ and the inner product $(\cdot, \cdot)$ are invariant under $\text{Ad}(N_G(H_0))$, where $N_G(H_0)$ is the normalizer of the connected unit component H_0 of H in G .*

Proof. Evidently, $\mathfrak{h}$ is $\text{Ad}(N_G(H_0))$ -invariant. Then $\mathfrak{p}$ is also $\text{Ad}(N_G(H_0))$ -invariant, because $\langle \cdot, \cdot \rangle$ is $\text{Ad}(G)$ -invariant. Now the $\text{Ad}(N_G(H_0))$ -invariance of $(\cdot, \cdot)$ follows from Theorem 9. ■

Remark 2. It follows from Theorems 8 and 30 that in general case the metric D on G is not Riemannian even in the case when (M, μ) is a δ -homogeneous Riemannian manifold. This is the reason for the words "in the generalized sense" in the statement of Theorem 4.

Theorem 10. *A Riemannian manifold (M, μ) is (G) - δ -homogeneous if and only if any of two following conditions are satisfied:*

1) *For every tangent vector $v \in M_x$, where x is any point in M , there is a Killing vector field X (in the Lie algebra RG of right-invariant vector fields on the Lie group G) on M such that $X(x) = v$ and $\mu(X(x), X(x)) = \max_{y \in M} \mu(X(y), X(y))$.*

2) *Every geodesic γ in M is an orbit of a 1-parameter motion group of M (in G) generated by a Killing vector field, attaining a maximal value of its length on γ .*

Proof. Let us remark at first that we can suggest that the vector v in the condition 1) is non-zero; then the condition 2) implies condition 1), while the condition 2) follows from the condition 1) and Proposition 5.7 of the chapter VI in [28], which states that an integral trajectory of a Killing vector field X on M , going through a point $x \in M$, is a geodesic, if x is a critical value of the function $\mu(X, X)$ and $X(x) \neq 0$.

Let suppose that (M, μ) is δ -homogeneous. Then Theorems 7 and 4 immediately imply the condition 2).

Sufficiency of 2). It's clear that the condition 2) implies that M is (G) -homogeneous. Then there is a constant $r > 0$ such that $\text{Radinj}(M) > r$. Let $x, y \in M$ and $\rho(x, y) = t < r$. Then there is unique geodesic $\gamma(s), s \in \mathbb{R}$, parameterized by arc length such that $\gamma(0) = x, \gamma(t) = y$. By the condition, $\gamma(s) = g(s)(x)$, where $g(s), s \in \mathbb{R}$, is a 1-parameter motion group of M (in G), generated by a Killing vector field X , such that $\mu(X(x), X(x)) = \max_{z \in M} \mu(X(z), X(z))$. Then it is clear that for every $z \in X$, $\rho(x, y) = \rho(x, g(t)(x)) \geq \rho(z, g(t)(z))$. We proved that M is restrictively (G) - δ -homogeneous. Hence M is (G) - δ -homogeneous by Proposition 2. ■

Definition 6. A Riemannian manifold (M, μ) is called *(G) -geodesic orbit* ((G) -g.o.), if every geodesic in M is an orbit of a one-parameter isometry subgroup (in G).

More extensive information on geodesic orbit manifolds (or *geodesic orbit spaces* by another terminology) one can find e.g. in [4, 31, 42, 43, 52].

Corollary 6. *Every (G) - δ -homogeneous Riemannian manifold is (G) -geodesic orbit ((G) -g.o.) manifold.*

4. TOTALLY GEODESIC SUBMANIFOLDS

In this section we investigate some totally geodesic submanifolds of δ -homogeneous and g.o. Riemannian manifolds.

Proposition 4 (Theorem 8.9 of Chapter VII in [28]). *Let M be a Riemannian manifold, N is its totally geodesic submanifold, X is a Killing field on M . Consider a smooth vector field $\tilde{X}$ on N , with is tangent (with respect to N) component of the field X . Then $\tilde{X}$ is a Killing field on the Riemannian manifold N .*

In [28] this proposition is used to prove that every closed totally geodesic submanifold of a homogeneous Riemannian manifold is homogeneous itself (Corollary 8.10 of Chapter VII in [28]). Here we give some refinement of this classical result.

Theorem 11. *Every closed totally geodesic submanifold of a δ -homogeneous Riemannian manifold is δ -homogeneous itself.*

Proof. Let N be a closed totally geodesic submanifold of a δ -homogeneous Riemannian manifold M . Since M is homogeneous, it is complete. Since N is closed submanifold of M , it is complete too. Let $U \neq 0$ be a tangent vector at some point $x \in N$. By Theorem 10 to prove the δ -homogeneity of N it is enough to show that there is a Killing field Y on N , whose value at the point x is U , and the maximal value of the length of Y is attained at the point x .

Since M is δ -homogeneous Riemannian manifold, there is a Killing field X on M such that its value at the point x is U , and the maximal value of its length is attained at the point x . Now as a required Killing field Y we can take $\tilde{X}$, the tangent component of the field X to N . According to Proposition 4, this field is Killing on N and $\tilde{X}(x) = X(x)$ obviously. Since at the point x the length of the field X is maximal among all points $y \in M$, then x is a point of maximal value for the length of the field $\tilde{X}$ (the length of the field $\tilde{X}$ does not exceed the length of the field X at all points of the manifold N). Theorem is proved. ■

Corollary 7. *Every closed totally geodesic submanifold of a normal homogeneous Riemannian manifold is δ -homogeneous.*

Remark 3. Let M be a Riemannian manifold, F is some set of its isometries. Then every connected component of the set of points of M , which are fixed under every isometry in F , is a closed totally geodesic submanifold of M . By the same manner, if K is some set of Killing fields on M , then every connected component of the set of points of M , which are zeros for every Killing field in K , is a closed totally geodesic submanifold of M [28].

Theorem 12. *Every closed totally geodesic submanifold of geodesic orbit (g.o.) Riemannian manifold is geodesic orbit itself.*

Proof. Let N be a closed totally geodesic submanifold of a geodesic orbit Riemannian manifold M . It is clear that M and N are complete. Let $U \neq 0$ be a tangent vector at some point $x \in \widetilde{M}$. It is enough to prove that there is a Killing field Y on N with the following properties:

- 1) the value Y at the point x is U ;
- 2) x is a critical point of the length of the field Y on N .

Indeed, in this case a geodesic passing through x in the direction U is an orbit of an one-dimensional motion group generated by the Killing field Y (this one-parameter group is correctly defined because of the completeness of N).

Since M is a geodesic orbit Riemannian manifold, there is a Killing field X on M , whose value at the point x is U , and such that x is a critical point of the length of the field X . Now as a required Killing field Y one can consider $\tilde{X}$, the tangent component of the field X to N . According to Proposition 4, it is a Killing field on N , and, moreover, $\tilde{X}(x) = X(x)$.

Now we need to prove only that x is a critical point of the length of the field $\tilde{X}$ on N . Let $Z = X - \tilde{X}$ be the normal component of the field X on the manifold N , and let g be the metric tensor on M . It is clear that

$$g(\tilde{X}, \tilde{X}) = g(X, X) - g(Z, Z).$$

The point x is a zero point for $g(Z, Z)$, therefore, x is a point of the minimal value of $g(Z, Z)$ on N . Consequently, x is a critical point both to the function $g(X, X)$ and to the function $g(Z, Z)$ on the manifold N . But in this case x is a critical point for the function $g(\tilde{X}, \tilde{X})$ also. Therefore, x is a critical point of the length of the field $\tilde{X}$ (since $\tilde{X}(x) = U \neq 0$). Theorem is proved. ■

According to Lemma 2, the metric product of δ -homogeneous spaces is δ -homogeneous itself. In the Riemannian case we have the conversion to this statement:

Theorem 13. *Let $M = M_0 \times M_1 \times \cdots \times M_k$ be a direct metric decomposition of a δ -homogeneous (respectively, g.o.) Riemannian manifold M with the maximal Euclidean factor M_0 . Then all factors of this product are δ -homogeneous (respectively, g.o.). If M is δ -homogeneous, then M_i are compact for $i \neq 0$. Besides, an isometry $f = f_0 \times \cdots \times f_k$ of the manifold M , which is a product of δ -translations, is a δ -translation itself.*

Proof. Since every fiber of the product under consideration is a complete totally geodesic submanifold, then according to Theorem 11 (Theorem 12), all factors are δ -homogeneous (respectively, g.o.), which proves the first statement. The second statement follows from the maximality of the Euclidean factor M_0 , Proposition 1 and Theorem 2. The last statement of Theorem follows from Lemma 2. ■

Since every g.o. (in particular, every δ -homogeneous) Riemannian manifold is homogeneous, it is useful to remind an algebraic description of homogeneous Riemannian manifolds. Let (M, μ) be a homogeneous Riemannian manifold with a closed connected transitive isometry group G , and H is its isotropy subgroup at a given point $x \in M$. Then M is naturally identified with the coset space G/H . Consider the Lie algebras $\mathfrak{h}$ and $\mathfrak{g}$, $\mathfrak{h} \subset \mathfrak{g}$, of the groups G and H . It is possible to choose some $\text{Ad}(H)$ -invariant complement $\mathfrak{p}$ to $\mathfrak{h}$ in $\mathfrak{g}$, which could be identified with the tangent space M_x of (M, μ) at the point x . In this case the homogeneous Riemannian metric μ is identified with some $\text{Ad}(H)$ -invariant inner product $(\cdot, \cdot)$ on $\mathfrak{p}$, whereas $\mathfrak{g}$ is identified with the Lie algebra of Killing vector fields on (M, μ) (see details in [12], Chapter VII).

Remark 4. If M is compact, then G is compact too, therefore, there exists some $\text{Ad}(G)$ -invariant inner product $\langle \cdot, \cdot \rangle$ on the Lie algebra $\mathfrak{g}$ of the group G . In this case as $\mathfrak{p}$ we can consider a $\langle \cdot, \cdot \rangle$ -orthogonal complement to $\mathfrak{h}$ in $\mathfrak{g}$. Note also that restrictions of $(\cdot, \cdot)$ and $\langle \cdot, \cdot \rangle$ to any $\text{Ad}(H)$ -invariant and $\text{Ad}(H)$ -irreducible submodule $\mathfrak{q} \subset \mathfrak{p}$ are proportional one to another.

Let $(M = G/H, \mu)$ be a homogeneous Riemannian manifolds with a closed connected transitive isometry group G , which is generated by some $\text{Ad}(H)$ -invariant inner product $(\cdot, \cdot)$ on $\mathfrak{p}$ in the above notation. For Killing fields $X, Y \in \mathfrak{p}$ we have the following equality:

$$\nabla_X Y(x) = -\frac{1}{2}[X, Y]_{\mathfrak{p}} + U(X, Y), \quad (4.4)$$

where the (bilinear symmetric) map $U : \mathfrak{p} \times \mathfrak{p} \rightarrow \mathfrak{p}$ is defined by the formula

$$2(U(X, Y), Z) = ([Z, X]_{\mathfrak{p}}, Y) + (X, [Z, Y]_{\mathfrak{p}}) \quad (4.5)$$

for any $Z \in \mathfrak{p}$ [12]. In [3] it is proved the following (compare with [46], Theorem 4.1)

Proposition 5 ([3]). *Let $(M = G/H, \mu)$ be any homogeneous Riemannian manifold and T be any torus in H , $C(T)$ is its centralizer in G . Then the orbit $M_T = C(T)(x)$ is a totally geodesic submanifold of (M, μ) .*

Proof. It is easy to get that the Lie subalgebra $\mathfrak{l}$ of $C(T)$ in $\mathfrak{g}$ has the form $\mathfrak{l} = \mathfrak{k} \oplus \mathfrak{q}$, where $\mathfrak{q} = \{X \in \mathfrak{p} \mid [X, \mathfrak{t}] = 0\}$, $\mathfrak{k} = \{X \in \mathfrak{h} \mid [X, \mathfrak{t}] = 0\}$, $\mathfrak{t} \subset \mathfrak{k}$ is the Lie algebra of T .

According to (4.4), to prove Proposition we need to show that $U(X, Y) \in \mathfrak{q}$ for any $X, Y \in \mathfrak{q}$. Let $W \in \mathfrak{p}$, $Z \in \mathfrak{t}$, then

$$\begin{aligned} 2([Z, U(X, Y)], W) &= -2(U(X, Y), [Z, W]) = -([Z, W], X)_{\mathfrak{p}}(Y) - (X, [Z, W]_{\mathfrak{p}})_{\mathfrak{p}}(Y) = \\ &= ([W, X], Z)_{\mathfrak{p}}(Y) + (X, [W, Y]_{\mathfrak{p}})_{\mathfrak{p}}(Z) = ([W, X]_{\mathfrak{p}}, [Z, Y]) + ([Z, X], [W, Y]_{\mathfrak{p}}) = 0. \end{aligned}$$

Since $W \in \mathfrak{p}$ may be chosen arbitrary, we have $[Z, U(X, Y)] = 0$ for any $Z \in \mathfrak{t}$. This means that $U(X, Y) \in \mathfrak{q}$. ■

Remark 5. If T is a maximal torus in H , then subalgebra $\mathfrak{k} = \mathfrak{t}$ is a part of the center of Lie algebra $\mathfrak{l}$. Therefore, in this case $\mathfrak{q}$ is the Lie algebra of some subgroup $Q \subset G$. Moreover, we can consider M_T as an orbit of Q through the point $x \in M$.

Now we consider some properties of g.o. manifolds. If we represent a homogeneous Riemannian metric μ on $M = G/H$ as a suitable $\text{Ad}(H)$ -invariant inner product $(\cdot, \cdot)$ on $\mathfrak{p}$ in the above notation, we can consider a useful notion of *geodesic vectors* on (M, μ) . A vector $X + Y$, where $Y \in \mathfrak{p}$ and $Y \in \mathfrak{h}$, is called *geodesic*, if the orbit of one-parameter group generated by the Killing field $X + Y$ is a geodesic of (M, g) , passing through the point $x \in M$ with stabilizer group H in the direction X . It is clear that a homogeneous Riemannian manifold $(G/H = M, \mu)$ is G-g.o. manifold if and only if for any $X \in \mathfrak{p}$ there is $Y \in \mathfrak{h}$ such that the vector $X + Y$ is geodesic. It is well known the following criterion for geodesic vectors (see e.g. [31]).

Proposition 6. *A vector $X + Y$, where $X \in \mathfrak{p}$ and $Y \in \mathfrak{h}$, is geodesic if and only if for every $V \in \mathfrak{p}$ the equality $([X + Y, V]_{\mathfrak{p}}, X) = 0$ holds.*

Proposition 7. *Let $(G/H, \mu)$ be a G-g.o.-space. For any $X \in \mathfrak{p}$ and $Y \in \mathfrak{h}$ such that $X + Y$ is geodesic vector we have the equality $U(X, X) = [X, Y]$, where U is defined by (4.5).*

Proof. For the geodesic vector $X + Y$ we have the equality

$$\begin{aligned} 0 &= (X, [V, X + Y]_{\mathfrak{p}}) = (X, [V, X]_{\mathfrak{p}}) + (X, [V, Y]) = (X, [V, X]_{\mathfrak{p}}) + ([Y, X], V) = \\ &= (U(X, X) + [Y, X], V) \end{aligned}$$

for every $V \in \mathfrak{p}$. Therefore, $U(X, X) = [X, Y]$. ■

Definition 7. A homogeneous Riemannian manifold (M, μ) is called (G) -*naturally reductive*, if there exist a connected Lie subgroup $G \subset \text{Isom}(M)$, acting transitively and effectively on M and a $\text{Ad}(H)$ -invariant decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$, where $\mathfrak{h}$ is the Lie algebra of the isotropy subgroup $H \subset G$ at some point in $x \in M$, such that one of the following equivalent statements holds:

- (1) every geodesic in M through the point x is an orbit of a one-parameter subgroup in G , generated by some $X \in \mathfrak{p}$;
- (2) $\mu([Z, X]_{\mathfrak{p}}, Y) + \mu(X, [Z, Y]_{\mathfrak{p}}) = 0$ for all $X, Y, Z \in \mathfrak{p}$ (in other words, $U \equiv 0$).

We obviously get from Proposition 7

Corollary 8. *Let $(G/H, \mu)$ be a G -g.o.-space. If for any $X \in \mathfrak{p}$ there is some $Z \in \mathfrak{h}$ such that $X + Z$ is geodesic vector and $[Z, X] = 0$, then $(G/H, \mu)$ is G -naturally reductive.*

Now we get some simple general remarks.

Proposition 8. *Let $(G/H, \mu)$ be a G -g.o.-space. Consider any $\text{Ad}(H)$ -invariant submodule $\mathfrak{q} \subset \mathfrak{p}$. Then for every $X, Y \in \mathfrak{q}$ we have $U(X, Y) \in \mathfrak{q}$.*

Proof. Consider some geodesic vectors $X + Z_1, Y + Z_2, X + Y + Z_3$, where $X, Y \in \mathfrak{q}$ and $Z_i \in \mathfrak{h}$. We get from Proposition 7 that $U(X, X) = [X, Z_1] \in \mathfrak{q}$, $U(Y, Y) = [Y, Z_2] \in \mathfrak{q}$, $U(X + Y, X + Y) = [X + Y, Z_3] \in \mathfrak{q}$. Therefore, $2U(X, Y) = U(X + Y, X + Y) - U(X, X) - U(Y, Y) \in \mathfrak{q}$. ■

Proposition 9. *Let $(G/H, \mu)$ be a G -g.o. manifold (G - δ -homogeneous manifold), and L is a Lie subgroup of G such that $H \subset L \subset G$. Then the orbit of the group L through the point x in G/H is a totally geodesic submanifold of $(G/H, \mu)$. In particular, L/H with the metric, induced by μ , is g.o. space (respectively, δ -homogeneous space).*

Proof. Let $\mathfrak{l}$ be a Lie algebra of L . Consider the decomposition $\mathfrak{l} = \mathfrak{h} \oplus \mathfrak{q}$, where $\mathfrak{q} = \mathfrak{p} \cap \mathfrak{l}$. Then the module $\mathfrak{q} \subset \mathfrak{p}$ is $\text{Ad}(H)$ -invariant. According to Proposition 8 we have $U(X, Y) \in \mathfrak{q}$ for every $X, Y \in \mathfrak{q}$. On the other hand, for every $X, Y \in \mathfrak{q}$ we have $[X, Y] \in \mathfrak{l} = \mathfrak{h} \oplus \mathfrak{q}$. Therefore, by (4.4) we get $\nabla_X Y(x) \in \mathfrak{q}$ for any $X, Y \in \mathfrak{q}$. This means that the homogeneous submanifold L/H (with the induced metric) is totally geodesic in $(G/H, \mu)$. The last statement follows from Theorem 12 (respectively, 11). ■

At the end of this section we note one special property of compact homogeneous Riemannian manifolds.

Proposition 10. *Let $(M = G/H, \mu)$ be a compact homogeneous Riemannian manifold. Consider any $\text{Ad}(H)$ -invariant and $\text{Ad}(H)$ -irreducible submodule $\mathfrak{q} \subset \mathfrak{p}$, where $\mathfrak{p}$ is a $\langle \cdot, \cdot \rangle$ -orthogonal complement to $\mathfrak{h}$, and $\langle \cdot, \cdot \rangle$ is some $\text{Ad}(G)$ -invariant inner product on $\mathfrak{g}$ (see Remark 4). Then for every $X, Y \in \mathfrak{q}$ we have $U(X, Y) = 0$.*

Proof. Since the module $\mathfrak{q}$ is $\text{Ad}(H)$ -invariant and $\text{Ad}(H)$ -irreducible, $(\cdot, \cdot)|_{\mathfrak{q}} = \alpha \langle \cdot, \cdot \rangle|_{\mathfrak{q}}$ for some $\alpha > 0$. Therefore, for any $Z \in \mathfrak{p}$ we have

$$2(U(X, Y), Z) = ([Z, X]_{\mathfrak{p}}, Y) + (X, [Z, Y]_{\mathfrak{p}}) = \alpha([Z, X]_{\mathfrak{p}}, Y) + \alpha(X, [Z, Y]_{\mathfrak{p}}) = 0,$$

since $\langle \cdot, \cdot \rangle$ is $\text{Ad}(G)$ -invariant. ■

5. ADDITIONAL SYMMETRIES OF δ -HOMOGENEOUS METRICS

Remind that the group G acts on the homogeneous space G/H by the transformation $L_b : G/H \rightarrow G/H$ ($b \in G$), where

$$L_b(cH) = bcH.$$

Let $N_G(H)$ be the normalizer of H in the group G . For every $a \in N_G(H)$ one can correctly define a G -equivariant diffeomorphism $R_a : G/H \rightarrow G/H$ acting by the following rule:

$$R_a(cH) = cHa^{-1} = ca^{-1}H.$$

Theorem 14. *Let $(G/H, \rho)$ be a compact G - δ -homogeneous Riemannian manifold with a connected transitive isometry group G , $N_G(H)$ is the normalizer of the subgroup H in the group G . Then for every $a \in N_G(H)$ the diffeomorphism $R_a : G/H \rightarrow G/H$ is a Clifford-Wolf translation on the Riemannian manifold $(G/H, \rho)$.*

Proof. It is clear that the isometricity of the map R_a is equivalent to that that for all elements $c \in G$ and $a \in N_G(H)$, the differential $dr_{a^{-1}}(c)$ preserves the length of every vector $u \in \text{hor}_c \subset G_c$, where hor_c means the horizontal subspace of the corresponding Riemannian submersion $pr : (G, \nu) \rightarrow (G/H, \mu)$ in G_c and $dr_{a^{-1}}(\text{hor}_c) = \text{hor}_{ca^{-1}}$. Here r, l denote the operations of right and left translations in G . We have the evident equality

$$r_{a^{-1}} = l_c \circ l_{a^{-1}} \circ (l_a \circ r_{a^{-1}}) \circ l_{c^{-1}},$$

and the corresponding composition of their differentials. Now it is clear that $l_{c^{-1}}(c) = e$, $dl_{c^{-1}}(\text{hor}_c) = \text{hor}_e = p$, and $d(l_a \circ r_{a^{-1}})(e) = \text{Ad}(a)$. But the last map preserves the space p and the scalar product $(\cdot, \cdot)$ by Corollary 5 and evident inclusion $N_G H \subset N_G(H_0)$. All differentials of left translations preserve the horizontal distribution and length of horizontal vectors. So, the map R_a is an isometry. It is a Clifford-Wolf translation, because it is generated by the right translation r_a of G , commuting with all left translations of G , which generate a transitive isometry group of $(G/H, \rho)$. ■

Lemma 3. *The transformation R_a of (effective) homogeneous space G/H for $a \in N_G(H)$ coincides with a transformation L_b for some $b \in G$ if and only if a is the product of some central element of the group G and some element of the group H .*

Proof. Suppose that $R_a = L_b$ for some $b \in G$. Since R_a obviously commutes with every transformation L_d , $d \in G$, we obtain that b is in the center of G . Further, the condition $R_a = L_b$ is equivalent to the next one: $ca^{-1}H = bcH = cbH$ for any $c \in G$. Therefore, $a = \tilde{b}d$, where $\tilde{b} = b^{-1}$ is a central element of G , and d is some element of the group H . The converse is obvious. ■

Theorem 15. *Let $(G/H, \rho)$ be a compact δ -homogeneous Riemannian manifold with a closed connected transitive semisimple isometry Lie group G . Then the group $N_G(H)/H$ is finite.*

Proof. According to Theorem 14, for every $a \in N_G(H)$ the diffeomorphism $R_a := G/H \rightarrow G/H$, acting by the rule $R_a(cH) = cHa^{-1} = ca^{-1}H$, is an isometry of $(G/H, \rho)$.

If $\dim(N_G(H)) > \dim(H)$, then one can choose a continuous family of isometries of the form R_a , which are not in the group G . Really, let us consider a vector U , which is in the Lie algebra of the group $N_G(H)$, but not in $\mathfrak{h}$. Consider $a = \exp(tU) \in N_G(H)$ for some real number t . Then the transformation R_a is an isometry of $(G/H, \rho)$. Since the center of the group G is discrete, with using of Lemma 3 we get that for some open set $O \subset \mathbb{R}$ all the transformations R_a for $a \in O$ are not in the group G . But this contradicts to the fact that G is the full connected isometry group of the Riemannian manifold $(G/H, \rho)$.

Therefore, we conclude that $\dim(N_G(H)) = \dim(H)$, and the group $N_G(H)/H$ is finite, since it is compact. ■

Example 3. Let G be a connected compact semisimple Lie group, and μ is some left-invariant Riemannian metric on G , so G is a closed connected transitive isometry group of the Riemannian manifold (G, μ) . Then (G, μ) is not G - δ -homogeneous. Really, if (G, μ) is G - δ -homogeneous, then according to Theorem 15, the group $N_G(H)/H$ is finite. But in our case $H = \{e\}$ is trivial, and $N_G(H)/H = G$ is not discrete.

According to the previous example we need to discuss δ -homogeneous left-invariant metrics on compact Lie groups. It is clear that any bi-invariant metric ρ on a compact Lie group G is G - δ -homogeneous. But there exist δ -homogeneous left-invariant metrics on G which are not bi-invariant. One can show this as follows.

Let G be a compact connected semisimple Lie group, and let K be a connected subgroup of G . Among all left-invariant metrics on G we consider a subclass $\mathcal{M}_{G,K}$ of metrics which

are right-invariant with respect to K . It is easy to see that the subclass $\mathcal{M}_{G,K}$ consists of $(G \times K)$ -invariant metrics on the homogeneous space $M = (G \times K)/\text{diag}(K)$ (we use the natural inclusion $K \subset G$). Indeed, every metric from $\mathcal{M}_{G,K}$ has $G \times K$ as a transitive motion group with the isotropy subgroup $\text{diag}(K)$ at the unit $e \in G$. On the other hand, it is clear that G is transitive on the space $M = (G \times K)/\text{diag}(K)$.

Now let us consider a $(G \times K)$ -normal homogeneous metric ρ on M . Then the Riemannian homogeneous space (M, ρ) is $(G \times K)$ - δ -homogeneous (Corollary 1). But the above discussion implies that (M, ρ) is isometric to the Lie group G with some left-invariant metric ρ_1 . This metric could be bi-invariant, but it is easy to see that the set of $(G \times K)$ -normal homogeneous metric ρ on M is more extensive than the set of bi-invariant metrics on G (for more details see [19]). Therefore, we obtain δ -homogeneous left-invariant metrics on G which are not bi-invariant.

Example 4. Let F be a connected compact simple Lie group, $G = F^k$, $k \geq 2$, $H = \text{diag}(F) \subset G$. Let us consider the space $G/H = F^k/\text{diag}(F)$ supplied with the metrics ρ generated by the Killing form of F^k . Then the homogeneous Riemannian manifold $(G/H, \rho)$ is δ -homogeneous. On the other hand it is isometric to the Lie group F^{k-1} with some left-invariant metric ρ_1 . If $k \geq 3$, the metric ρ_1 is not bi-invariant.

Remark 6. It is obvious that for a compact G - δ -homogeneous Riemannian manifold $(G/H, \rho)$ with positive Euler characteristic all conditions of Theorem 15 are fulfilled. Really, any connected one-dimensional central subgroup of G induces on G/H a non-vanishing vector field, but this implies that $\chi(G/H) = 0$. On the other hand, in the case of positive Euler characteristic the statement of Theorem 15 is well known, since the groups H and G have one and the same rank.

6. δ -VECTORS

Let suppose that $M = (G/H, \mu)$ be a compact homogeneous connected Riemannian manifold with connected (compact) Lie group G . Let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$, $\langle \cdot, \cdot \rangle$, and $(\cdot, \cdot)$ be the same as in the section 4. We identify the Lie algebra of Killing vector fields on M with the Lie algebra $\mathfrak{g}$ of right invariant vector fields on G and use $\text{Ad}(G)$ -invariant (Chebyshev's) norm $\|\cdot\|$ on $\mathfrak{g}$ and corresponding bi-invariant inner metric D on G from Theorem 6.

From Section 3 we get the following

Proposition 11. *The map $p : (G, D) \rightarrow (G/H, \mu)$ does not increase distances. It is a submetry if and only if M is G - δ -homogeneous.*

Definition 8. A vector $w \in \mathfrak{g}$ is called δ -vector on the Riemannian homogeneous manifold $(M = G/H, \mu)$ if $|P(w)| := \sqrt{(P(w), P(w))} = \|w\|$, where P is as in Theorem 9. (This is equivalent to the condition that for any $a \in G$, $(w_{\mathfrak{p}}, w_{\mathfrak{p}}) \geq (\text{Ad}(a)(w)|_{\mathfrak{p}}, \text{Ad}(a)(w)|_{\mathfrak{p}})$.)

Proposition 12. *Let suppose that for a vector $v \in \mathfrak{p}$, the set $W(v)$ of all δ -vectors of the form $w = v + u$, $u \in \mathfrak{h}$ (such that $\|w\| = \sqrt{(v, v)}$) is non empty. Then $W(v)$ is compact and convex. Moreover, there is a unique vector $w = w(v) \in W(v)$ with the smallest distance $\sqrt{\langle w - v, w - v \rangle}$.*

Proof. We can suggest that $\sqrt{(v, v)} = 1$. Since p in Proposition 11 doesn't increase distances, then P in Theorem 9 has the same property, and really $\|w\| = 1$. Let suppose that $w_1, w_2 \in W(v)$, $0 \leq t \leq 1$, and $w = tw_1 + (1-t)w_2$. Then by the triangle inequality,

$$\|w\| = \|tw_1 + (1-t)w_2\| \leq t\|w_1\| + (1-t)\|w_2\| = t + (1-t) = 1.$$

Since P is a linear map, then

$$P(w) = P(tw_1 + (1-t)w_2) = tP(w_1) + (1-t)P(w_2) = tv + (1-t)v = v.$$

One more, because P doesn't increase distances, it follows from the last two relations that $\|w\| = 1$ and $w \in W(v)$. So, the set $W(v)$ is convex. Evidently, it is compact, and we proved the first statement.

It follows from compactness of $W(v)$ the existence of a vector $w \in W(v)$ with the smallest $|w-v|_1 = \sqrt{\langle w-v, w-v \rangle}$. If we have another such a vector $w' \neq w$, then by the previous statement, $w'' := \frac{1}{2}(w+w') \in W(v)$ and

$$2|w''-v|_1 = |(w-v) + (w'-v)|_1 < |(w-v)|_1 + |(w'-v)|_1 = 2|w-v|_1,$$

a contradiction. ■

Remark 7. Let $v \in \mathfrak{p}$ with $W(v) \neq \emptyset$. According to Proposition 12, there is a unique vector $w \in W(v)$ with the smallest distance $\sqrt{\langle w-v, w-v \rangle}$. Later on we shall use a notation $w(v)$ for this vector and a notation $u(v)$ for the vector $w(v) - v \in \mathfrak{h}$.

Proposition 13. Consider any vector $v \in \mathfrak{p}$ with $W(v) \neq \emptyset$. The following four statements are equivalent (see Remark 7): $w(v) = v$, $u(v) = 0$, $\|v\| = |v|$, and the corresponding vector field $X(v)$ on M is infinitesimal $\delta(x_0)$ -translation for the point $x_0 = p(e)$.

Proposition 14. If $W(v) \neq \emptyset$, then the inequalities $u(v) \neq 0$ and $\|v\| > |v|$ are equivalent. In this case the following statements are satisfied: for every element $g \in G$, such that $\text{Ad}(g)(\mathfrak{h}) = \mathfrak{h}$, the equality $\text{Ad}(g)(v) = v$ (respectively, $\text{Ad}(g)(v) = -v$) implies that $\text{Ad}(g)(u(v)) = u(v)$ (respectively, $\text{Ad}(g)(u(v)) = -u(v)$).

Proof. This follows easily from Propositions 12, 11 and the fact that $\|\cdot\|$, $\langle \cdot, \cdot \rangle_e$ are $\text{Ad}(G)$ -invariant and invariant under central symmetry. ■

From Theorem 9 we get the following

Proposition 15. A homogeneous Riemannian manifold $(G/H, \mu)$ with connected Lie group G is G - δ -homogeneous if and only if for every vector $v \in \mathfrak{p}$ there exists a vector $u \in \mathfrak{h}$ such that the vector $v + u$ is a δ -vector.

7. ON THE TOPOLOGY OF COMPACT HOMOGENEOUS SPACES

In general case a Cartan subalgebra $\mathfrak{k}$ of a Lie algebra $\mathfrak{g}$ is defined as a nilpotent Lie subalgebra in $\mathfrak{g}$, which coincides with its normalizer in $\mathfrak{g}$. If a Lie algebra $\mathfrak{g}$ is compact, i.e. is the Lie algebra of some compact Lie group G , then $\mathfrak{k}$ is a maximal commutative subalgebra in $\mathfrak{g}$, hence, is the Lie algebra of a maximal torus T in G .

Theorem 16 ([1]). Any two maximal tori in a compact (connected) Lie group G are conjugate by an inner automorphism of the Lie group G .

Thus, the rank $\text{rk}(G)$ of a compact Lie group G is (correctly) defined as the dimension of a Cartan subalgebra $\mathfrak{k}$ in $\mathfrak{g}$, or, what is equivalent, the dimension of a maximal torus in G .

Theorem 17 ([24], [37]). Let $M = G/H$ be a homogeneous space, where G, H are connected compact Lie groups. Then $\chi(M) \geq 0$. The following statements are equivalent: (i) $\chi(M) > 0$; (ii) $\text{rk}(G) = \text{rk}(H)$. If $\chi(M) > 0$, then the manifold M is formal and $\chi(M) = \frac{|W_G|}{|W_H|}$, where $|W_G|$ (respectively, $|W_H|$) is the order of the Weyl group W_G (respectively, W_H) of the Lie group G (respectively, H).

Theorem 18. Let $M = (G/H, \mu)$ be a compact simply connected homogeneous Riemannian manifold with compact Lie groups G and H , and G is connected. Then the following conditions are equivalent:

- 1) $\chi(M) = 0$;
- 2) $\text{rk } G > \text{rk } H$;
- 3) There is a right-invariant vector field on G , projecting under canonical map $p : G \rightarrow M$ to nowhere vanishing Killing field on M ;
- 4) All characteristic numbers of the Riemannian manifold M , defined for principal bundle $\pi : SO(M) \rightarrow M$ of orthonormal oriented bases on M , are equal to zero.

Proof. It follows from homotopic sequence of the bundle $p : G \rightarrow G/H$, connectedness of G , and simply connectedness of G/H that the group H is connected. So, all conditions of Theorem 17 are satisfied. Then the conditions 1) are 2) equivalent.

It is clear that the condition 3) implies the condition 1).

We will show that the condition 2) implies the statement 3).

Let us consider $U \in \mathfrak{g}$ such that the dimension of the closure in G of one-parameter group $\exp(tU)$ coincides with $\text{rk}(G)$, which, in turn, is strongly greater than $\text{rk}(H)$. We state that $\text{Ad}(s)(U) \notin \mathfrak{h}$ for all $s \in G$. Actually, let suppose that $V := \text{Ad}(s)(U) \in \mathfrak{h}$. Since $\text{Ad}(s)$ is an inner automorphism of Lie algebra $\mathfrak{g}$, then the dimension of the closure in G of one-parameter group $\exp(tV)$ also coincides with $\text{rk}(G)$. On the other hand, this closure is a torus in H , because H is a closed Lie subgroup of Lie group G . This contradicts to the inequality $\text{rk}(H) < \text{rk}(G)$. Clearly, a right-invariant vector field W on G with the condition $W(e) = U$ projects under the map p to a Killing vector field on M without zeroes.

Since any two maximal tori in a compact Lie group are conjugate, then one can easily prove that the condition 3) implies the condition 2), because the equality $\text{rk}(G) = \text{rk}(H)$ implies that every maximal torus is conjugate by an inner automorphism of Lie group G to a subgroup in H . Thus every right-invariant vector field on G projects to a Killing field on M , which necessarily vanishes at some points.

Characteristic numbers from the condition 4) are defined only for even-dimensional Riemannian manifold M . In this case also Euler characteristic is a characteristic number (corresponding to the characteristic Euler class) by Gauss-Bonnet theorem. Then in this case the condition 1) follows from the condition 4); The statement 4) follows from the condition 3) (even from the more weaker existence condition of nowhere vanishing Killing vector field on arbitrary compact smooth oriented Riemannian manifold of even dimension) by Bott's theorem [15] (a proof is also given in Theorem 6.1 of Chapter 2 in [27]).

In odd-dimensional case $\chi(M) = 0$ and the condition 1) is satisfied, hence 2) and 3), as we said before. If we suggest that characteristic numbers of odd-dimensional (compact Riemannian) manifold are equal zero by definition, then the condition 4) is automatically satisfied. Thus, in this case all 4 conditions are equivalent and always satisfied. ■

Proposition 16 ([46]). *Every even-dimensional homogeneous Riemannian manifold M of positive sectional curvature has positive Euler characteristic.*

Proof. According to Berger's theorem [11], any Killing field on an even-dimensional Riemannian manifold of positive sectional curvature must vanish at some point. If $M = G/H$ would have zero Euler characteristic, then by Theorem 18 M would admit nowhere vanishing Killing vector field. Thus $\chi(M) > 0$ by Hopf-Samelson theorem. ■

Remark 8. Example of a flat even-dimensional torus, which has zero Euler characteristic, shows that the statement of Proposition 16 is not true under the condition of non-negativeness of sectional curvature. Notice that by Poincaré duality, any compact odd-dimensional triangulated (in particular, smooth) manifold has zero Euler characteristic.

Corollary 9. *All CROSS'es, besides odd-dimensional, i.e. besides S^{2k+1} and $\mathbb{R}P^{2k+1}$, have positive Euler characteristic.*

Theorem 19. *Any simply connected compact homogeneous Riemannian manifold (M, g) admits a semi-simple compact transitive isometry group. If moreover the connected component of the group of all isometries of the space (M, g) is not semi-simple, then $\chi(M) = 0$ and (M, g) is a total space of a Riemannian submersion, which is a non-trivial principal bundle with simply connected homogeneous Riemannian base (M_1, g_1) and pair-wise isometric totally geodesic flat tori as fibers. Under this the connected component of the group of all motions of the space (M_1, g_1) is semi-simple. If (M, g) is δ -homogeneous, then (M_1, g_1) is also δ -homogeneous.*

Proof. The proof follows the line of the paper [7]. The first statement of theorem we get on the ground of Corollary 4 of the section 3 in the chapter 2 in [22].

Under this the connected component G of the full isometry group of the space (M, g) is not semi-simple if and only if G has non-trivial connected component C of it's center. Then the group C acts as a non-trivial connected group of Clifford-Wolf translations on (M, g) . Thus $\chi(M) = 0$.

It is clear that the orbits of one-parameter subgroups of the group C in (M, g) are geodesic (see also [7]). Thus the orbits of the group C are pair-wise isometric flat totally geodesic tori in (M, g) .

The simply connectedness of M and connectedness of fibers of Riemannian submersion $p : (M, g) \rightarrow (M_1, g_1)$ imply the non-triviality of the bundle p and simply connectedness of the space M_1 .

On the ground of Theorem 3, the metric quotient (orbit) space $(C \backslash M, g_1) := (M_1, g_1)$ is δ -homogeneous Riemannian manifold, if (M, g) is δ -homogeneous Riemannian manifold. ■

Remark 9. If (M, g) is a homogeneous compact Riemannian manifold and $\chi(M) > 0$, then by Theorem 19, the connected component (of effective) full isometry group of the manifold (M, g) is semi-simple. The opposite statement is not true: the connected component of full isometry group of Euclidean sphere S^{2l-1} , $l \geq 3$, is simple Lie group $SO(2l)$ and semi-simple Lie group $SO(4)$ with Lie algebra $so(4) = so(3) \oplus so(3)$ in the case of the sphere S^3 .

Remark 10. The well-known example of Berger spheres $S^{2n+1} = U(n+1)/U(n)$ shows that in general case the connected component G of the unit for full isometry Lie group of the space (M, μ) is not semi-simple, (even if (M, g) is normal); in this case the universal covering Lie group of G is non-compact. One needs to note also that for Berger spheres $U(n+1)/U(n)$ (with normal metrics) the Lie algebra of isotropy group $U(n)$ is not orthogonal to the center of Lie algebra $u(n+1)$ with respect to corresponding $\text{Ad}(U(n+1))$ -invariant scalar product.

It follows from Proposition 1 that every δ -homogeneous Riemannian manifold has non-negative sectional curvature.

Question 2. *Whether every compact δ -homogeneous Riemannian manifold with a finite fundamental group has positive Ricci curvature?*

Proposition 17. *Let $\mathfrak{h}$ is a Lie subalgebra of a Lie algebra $\mathfrak{g}$ of a connected Lie group G and $N_{\mathfrak{g}}(\mathfrak{h}) = \mathfrak{h}$, where $N_{\mathfrak{g}}(\mathfrak{h})$ is the normalizer of $\mathfrak{h}$ in $\mathfrak{g}$. Then $\mathfrak{h}$ is a Lie algebra of a unique closed connected Lie subgroup H in G .*

Proof. Let $H_1 = \{g \in G : \text{Ad}(g)(\mathfrak{h}) \subset \mathfrak{h}\}$. Then H_1 is closed subgroup in G . Hence its connected component H is closed. By Cartan theorem, H is a Lie subgroup of G . Evidently, Lie algebra of H is equal to $N_{\mathfrak{g}}(\mathfrak{h})$, which is by condition is equal to $\mathfrak{h}$, so H is required Lie subgroup. ■

One can easily deduce from this the following statements.

Proposition 18. *If $\mathfrak{h}$ is a reductive Lie subalgebra of $\mathfrak{g}$, containing a maximal commutative subalgebra $\mathfrak{t}$ in $\mathfrak{g}$, then $N_{\mathfrak{g}}(\mathfrak{h}) = \mathfrak{h}$.*

Theorem 20. *Let G be a simple compact connected Lie group and $\mathfrak{t}$ be Lie algebra of a maximal torus $T \subset G$. Then every proper Lie subalgebra $\mathfrak{h} \subset \mathfrak{g}$, such that $\mathfrak{t} \subset \mathfrak{h}$, is a Lie algebra of the unique closed connected Lie subgroup $H \subset G$. Moreover, G/H is a simply connected compact connected homogeneous space of positive Euler characteristic.*

8. HOMOGENEOUS SPACES OF POSITIVE EULER CHARACTERISTIC

Here we recall some properties of homogeneous spaces with positive Euler characteristic.

Theorem 21 ([41]). *If M and M' are homogeneous spaces of connected compact Lie groups, $\chi(M) > 0$, $\chi(M') > 0$ and M is homotopically equivalent to M' then M and M' are diffeomorphic.*

Now we outline some structure results about homogeneous spaces of positive Euler characteristic (see [37], 19.5). Let G/H be an almost effective compact homogeneous space of positive Euler characteristic with connected group G . From Theorem 18 we know that the center of G is discrete (hence, G is semi-simple), and that there is a maximal torus $T \subset G$ such that $T \subset H$. Since the center of G is contained in every maximal torus of G , we get the following

Proposition 19 ([47]). *If a compact connected Lie group G acts effectively on the space $M = G/H$ of positive Euler characteristic, then the center of G is trivial.*

Theorem 22 ([30]). *Let $(G/H, \mu)$ be a simply connected compact almost effective homogeneous Riemannian manifold of positive Euler characteristic. Then $(G/H, \mu)$ is indecomposable if and only if G is simple. In particular, a simple and a non-simple compact Lie groups can not both act transitively and effectively as a group of motions on a compact Riemannian manifold M with positive Euler characteristic.*

A. Borel and J. de Siebenthal obtained in [13] the classification of subgroups with maximal rank of compact Lie groups (see also Section 8.10 in [50]). This classification give us a description of compact homogeneous spaces with positive Euler characteristic. A complete description of homogeneous spaces of classical Lie groups with positive Euler characteristic have been obtained also by H.C. Wang in [47].

We will concern later with special cases of compact homogeneous manifolds of positive Euler characteristic, namely, the (generalized) flag manifolds. They can be described as orbits M of a compact connected Lie group G by the adjoint representation. In other words, $M = G/H$, where $H = Z_G(S)$ is the centralizer of a non-trivial torus $S \subset G$; the Lie group H is always connected. Under this orbits of regular elements in $\mathfrak{g}$ are called *(full) flag manifolds*.

The chapter 8 in [12] contains the following statements: simply connected compact homogeneous Kähler manifolds are exactly (generalized) flag manifolds. Any latter manifold (admitting a canonical Kähler-Einstein structure, unique in a sense) is a rational complex algebraic (hence complex projective) manifold. In a special case $G = Sp(l)$, the stabilizer sub-groups, whose center is 1-dimensional, are sub-groups $U(l-m) \times Sp(m)$. Among the corresponding orbits $M_{l-m}^{Sp(l)}$, the only ones for which the normal metric is Kähler (hence Kähler-symmetric) are $M_1^{Sp(l)}$, that is $\mathbb{CP}^{2l-1} = Sp(l)/U(1) \times Sp(l-1)$, and $M_l^{Sp(l)}$, isomorphic to $Sp(l)/U(l)$, which is the manifold of totally isotropic complex l -subspaces of $\mathbb{C}^{2l}$. The space $M_l^{SO(2l+1)} = SO(2l+1)/U(l)$ is the manifold of complex flags of type l .

Using chapter 15 in [37], we can add more. Any (compact generalized) flag manifold M , supplied with the above mentioned canonical Kähler-Einstein structure, is isomorphic to $\mathbb{G}/\mathbb{H}$, where $\mathbb{G}$ is a complex connected Lie group and $\mathbb{H}$ is a closed complex parabolic Lie subgroup in $\mathbb{G}$. We recall that a connected complex Lie subgroup of $\mathbb{G}$ is called *parabolic*, if it contains a Borel subgroup of $\mathbb{G}$. A *Borel subgroup* in $\mathbb{G}$ is any its maximal connected solvable complex Lie subgroup. Thus M is a so-called *flag homogeneous space*. Under this, the corresponding complex structure on M is induced by complex structure on $\mathbb{G}$. Any parabolic subgroup of $\mathbb{G}$ contains $\text{Rad}(\mathbb{G})$, a normal subgroup in $\mathbb{G}$. Hence M is a flag homogeneous space of semi-simple complex Lie group $\mathbb{G}_0 := \mathbb{G}/\text{Rad}(\mathbb{G})$. Under this $M = G_0/H_0$, where G_0 is any compact real form of $\mathbb{G}_0$ and $H_0 = G_0 \cap \mathbb{H}_0$ for $\mathbb{H}_0 = \mathbb{H}/\text{Rad}(\mathbb{G})$.

It is proved in Corollary 7.12, p. 301 in [34] that a maximal connected Lie subgroup H of maximal rank in a compact connected Lie group G is a connected component of the normalizer (=of the centralizer) of some element $g \in G$. On the ground of this Corollary and connected results, the Table 5.1 in [34] is given of all maximal connected compact subgroups H of maximal rank (more exactly, their Lie subalgebras) in a compact connected simple Lie groups G . In particular, G/H is an orbit of the above mentioned element $g \in G$ with respect to the action of the group $I(G)$ of all inner automorphisms of the Lie group G . A (generalized) flag manifolds also can be considered as such orbits, when $g \in G$ is taken in a

diffeomorphic image $\exp_G(U)$, where U is an open ball with the center $0 \in \mathfrak{g}$ with respect to an $\text{Ad}(G)$ -invariant Euclidean metric on $\mathfrak{g}$.

Theorem 23 ([40]). *Let G and G' be connected compact Lie groups, $H \subset G$ and $H' \subset G'$ their connected Lie subgroups of maximal rank, provided that the natural action of G and G' on $M = G/H$ and $M' = G'/H'$ are locally effective. Suppose that the graded rings $H(M, \mathbb{Z})$ and $H(M', \mathbb{Z})$ are isomorphic. Then*

(i) *If $M = M_1 \times \cdots \times M_s$ and $M' = M'_1 \times \cdots \times M'_t$ are the canonical decompositions of M and M' , then $s = t$ and M_k is diffeomorphic to M'_k after an appropriate permutation of the factors.*

(ii) *If G and G' are simple then either the pairs (G, H) and (G', H') are locally isomorphic or (up to transposition) they are locally isomorphic to the pairs of the following list:*

$$G = SU(2n) \ (n \geq 2), H = S(U(1) \times U(2n-1)); \\ G' = Sp(n), H' = U(1) \cdot Sp(n-1); \quad M = M' = \mathbb{C}P^{2n-1}.$$

$$G = SO(7), H = SO(6); \quad G' = G_2, H' = SU(3); \quad M = M' = S^6.$$

$$G = SO(7), H = SO(5) \times SO(2); \quad G' = G_2, H' = SU(2) \cdot SO(2); \quad M = M' = Gr_{7,2}^+.$$

$$G = SO(2n) \ (n \geq 4), H = U(n); \quad G' = SO(2n-1), H' = U(n-1); \quad M = M' = I^0 Gr_{2n,n}^C.$$

Theorem 23 implies easily the classification of transitive actions of connected compact Lie groups on simply connected homogeneous spaces of positive Euler characteristic.

Moreover, from results of [38] and [40] we have

Theorem 24. *Let $(G/H, \mu)$ be a simply connected Riemannian homogeneous manifold of positive Euler characteristic, and G is a simple connected Lie group. Then the full connected isometry group of $(G/H, \mu)$ is G/C (C is the center of G), excepting the cases when $(G/H, \mu)$ is one of the following manifolds:*

1) $G/H = Sp(n)/U(1) \cdot Sp(n-1)$ ($n \geq 2$), μ - symmetric (Fubini) metric on $\mathbb{C}P^{2n-1} = SU(2n)/S(U(1) \times U(2n-1))$;

2) $G/H = SO(2n-1)/U(n-1)$ ($n \geq 4$), μ - symmetric metric on $I^0 Gr_{2n,n}^C = SO(2n)/U(n)$;

3) $G/H = G_2/SU(2) \cdot SO(2)$, μ - symmetric metric on $Gr_{7,2}^+ = SO(7)/SO(5) \times SO(2)$;

4) $G/H = G_2/SU(3)$ (strongly isotropy irreducible), μ - arbitrary G -invariant metric.

In the first three cases the metric μ is not G -normal, in the last case μ is metric of constant curvature on $S^6 = SO(7)/SO(6)$.

Proof. Using Proposition 19 and Theorem 23, we easily get the main statements. We need only to show that in Cases 1), 2), and 3) the metric μ is not G -normal. It follows from results of [38]. Really, in that paper the author proved that the full connected isometry group of a simply connected G -normal homogeneous space $M = G/H$ of a connected simple compact Lie group G , is $G \cdot \text{Aut}_G(M)^0$ (a locally direct product), where

$$\text{Aut}_G(M) = \{f \in \text{Diff}(M) \mid f(gx) = gf(x), g \in G, x \in M\},$$

excepting the following cases: $G_2/SU(3) = S^6$, $Spin(7)/G_2 = S^7$, $Spin(8)/G_2 = S^7 \times S^7$. Only one of these spaces (namely, $G_2/SU(3) = S^6$) has positive Euler characteristic. Moreover, it is strongly isotropy irreducible. We need to note also that $\text{Aut}_G(M)^0$ is trivial for spaces $M = G/H$ of positive Euler characteristic (it is easy to see from Theorem 22). ■

Now we describe the sets of G -invariant metrics on the spaces G/H from items 1), 2), 3) of Theorem 24. Note, that each of these spaces is a (generalized) flag manifold. Note also, that G -invariant metrics on the space $G/H = G_2/SU(3)$ constitutes a one-dimensional family of pairwise homothetic metrics.

Example 5. It is known (see e.g. [51]) that the set of G -invariant metrics on $G/H = Sp(n)/U(1) \cdot Sp(n-1)$ ($n \geq 2$) is two-parametric. More exactly, let $\langle \cdot, \cdot \rangle$ be an $\text{Ad}(Sp(n))$ -invariant inner product on the Lie algebra $\mathfrak{g} = sp(n)$. In this case $\mathfrak{h} = u(1) \oplus sp(n-1) \subset \mathfrak{k} := sp(1) \oplus sp(n-1) \subset \mathfrak{g}$. Let us consider an $\langle \cdot, \cdot \rangle$ -orthogonal decomposition

$$sp(n) = \mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p} = \mathfrak{h} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2,$$

where $\mathfrak{h} \oplus \mathfrak{p}_2 = \mathfrak{k} = sp(1) \oplus sp(n-1)$. Then the modules $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are $\text{Ad}(H)$ -invariant, $\text{Ad}(H)$ -irreducible, and pairwise inequivalent with respect to $\text{Ad}(H)$. Therefore, any $Sp(n)$ -invariant metric on $G/H = Sp(n)/U(1) \cdot Sp(n-1)$ is generated by one of inner products on $\mathfrak{p}$ of the form

$$\langle \cdot, \cdot \rangle = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2}$$

for some positive x_1 and x_2 . Note, that the subset of $SU(2n)$ -invariant (symmetric) metrics on G/H consists of the metrics with the relation $x_2 = 2x_1$. In this case the full connected isometry group is a quotient-group of $SU(2n)$ by its center, and the metric μ is $SU(2n)$ -normal, and $(Sp(n)/U(1) \cdot Sp(n-1), \mu)$ is isometric to the complex projective space $\mathbb{C}P^{2n-1} = SU(2n)/U(1) \cdot S(U(2n-1))$ with the Fubini metric. Note also, that any $Sp(n)$ -invariant metric on $Sp(n)/U(1) \cdot Sp(n-1)$ is weakly symmetric and, hence, g.o.-metric [52].

Example 6. The set of G -invariant metrics on $G/H = SO(2n-1)/U(n-1)$ ($n \geq 3$) is two-parametric also. More exactly, let $\langle \cdot, \cdot \rangle$ be an $\text{Ad}(SO(2n-1))$ -invariant inner product on the Lie algebra $\mathfrak{g} = so(2n-1)$. In this case $\mathfrak{h} = u(n-1) \subset \mathfrak{k} := so(2n-2) \subset \mathfrak{g} = so(2n-1)$. Let us consider an $\langle \cdot, \cdot \rangle$ -orthogonal decomposition

$$so(2n-1) = \mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p} = \mathfrak{h} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2,$$

where $\mathfrak{h} \oplus \mathfrak{p}_2 = \mathfrak{k} = so(2n-2)$. Then the modules $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are $\text{Ad}(H)$ -invariant, $\text{Ad}(H)$ -irreducible, and pairwise inequivalent with respect to $\text{Ad}(H)$. Therefore, any $SO(2n-1)$ -invariant metric on $G/H = SO(2n-1)/U(n-1)$ is generated by one of inner products on $\mathfrak{p}$ of the form

$$\langle \cdot, \cdot \rangle = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2}$$

for some $x_1 > 0$ and $x_2 > 0$. Note, that the subset of $SO(2n)$ -invariant (symmetric) metrics on G/H consists of the metrics with the relation $x_2 = 2x_1$ [26]. As in the previous case, every $SO(2n-1)$ -invariant metric on $SO(2n-1)/U(n-1)$ is weakly symmetric and, hence, g.o.-metric [52]. Note also that $SO(5)/U(2)$ coincides with $Sp(2)/U(1) \cdot Sp(1)$ as a homogeneous space.

Example 7. Let us consider now the space $G/H = G_2/SU(2) \cdot SO(2)$, where $H = SU(2) \cdot SO(2) \subset SU(3)$, and $G_2/SU(3)$ is strongly isotropy irreducible ($G/H = SO(7)/SO(5) \times SO(2) = Gr_{7,2}^+$). It is easy to see that there is a subgroup $SO(4) \subset G_2$ such that $SU(2) \cdot SO(2) = SU(3) \cap SO(4)$. Therefore, we have $\langle \cdot, \cdot \rangle$ -orthogonal decomposition

$$g_2 = \mathfrak{h} \oplus \mathfrak{p} = \mathfrak{h} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3,$$

where $\langle \cdot, \cdot \rangle$ is some $\text{Ad}(G_2)$ -invariant inner product on g_2 , $su(3) = \mathfrak{h} \oplus \mathfrak{p}_3$, $so(4) = \mathfrak{h} \oplus \mathfrak{p}_2$, $\dim(\mathfrak{p}_2) = 2$, $\dim(\mathfrak{p}_1) = \dim(\mathfrak{p}_3) = 4$, and every module $\mathfrak{p}_i$ is $\text{Ad}(G_2)$ -invariant and $\text{Ad}(G_2)$ -irreducible. Moreover, the modules $\mathfrak{p}_1$, $\mathfrak{p}_2$, and $\mathfrak{p}_3$ are pairwise inequivalent with respect to $\text{Ad}(H)$ [26]. Therefore, we have 3-parametric family of G_2 -invariant metrics on G/H , every of each is generated by some inner product

$$\langle \cdot, \cdot \rangle = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2} + x_3 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_3}$$

on $\mathfrak{p}$ for some positive x_i , $i = 1, 2, 3$. From [26] we know that $SO(7)$ -invariant (symmetric) metrics on G/H are exactly metrics with the following relations:

$$x_2 = 2x_1, \quad x_3 = 3x_1.$$

Remind, that $G_2/SU(2) \cdot SO(2)$ is a flag manifold. The results of the paper [4] implies that any G_2 -invariant g.o.-metric μ on the space $G_2/SU(2) \cdot SO(2)$ is either G_2 -normal or $SO(7)$ -normal (symmetric).

Now we shall give a simple description of naturally reductive homogeneous manifolds of positive Euler characteristic, which follows from Theorem 22.

Theorem 25. *Let M be a compact naturally reductive homogeneous Riemannian manifold of positive Euler characteristic. Then M is G_1 -normal homogeneous for some (transitive on M) semi-simple Lie subgroup $G_1 \subset G$, where G is the full connected isometry group of M .*

Proof. The group G is semisimple, since $\chi(M) > 0$ (see Theorem 19).

In the proof of the statement of Theorem we can assume without loss of generality that M is simply connected. Really, the universal Riemannian covering $\widetilde{M}$ of M has a semisimple transitive group of motion $\widetilde{G}$, which is a covering of G . Since G and $\widetilde{G}$ have one and the same Lie algebra, $\widetilde{G}$ is compact, therefore, $\widetilde{M}$ is compact too. If $\widetilde{M}$ is normal homogeneous with respect to some semisimple subgroup $\widetilde{G}_1 \subset \widetilde{G}$, then M is G_1 -normal homogeneous, where $G_1 \subset G$ is the image of $\widetilde{G}_1$ under the natural covering epimorphism $\pi : \widetilde{G} \rightarrow G$.

Moreover, we can assume in addition that M is indecomposable. Really, if $M = M_1 \times \dots \times M_s$ is the de Rham decomposition of M then every M_i is naturally reductive homogeneous manifold ([29], Corollary 7; see also [28], Chapter X, theorem 5.2). If we prove that every M_i is normal homogeneous (with respect to some transitive subgroup of its full connected isometry group), then M is normal homogeneous too.

Let M be a compact simply connected indecomposable naturally reductive homogeneous manifold with $\chi(M) > 0$, and G is its (semisimple) connected isometry group. From Kostant theorem (Theorem 4 in [29]) we get that there is a subgroup $G_1 \subset G$, transitive on M , with the following property: there is an $\text{Ad}(G_1)$ -invariant non-degenerate quadratic form Q on the Lie algebra $\mathfrak{g}_1$ of the group G_1 such that the Riemannian metric of M is generated by the restriction of Q to Q -orthogonal complement $\mathfrak{p}$ to $\mathfrak{h}_1$ in $\mathfrak{g}_1$ (H_1 is the stabilizer group of some point of M with respect to the action of G_1 , and $\mathfrak{h}_1$ is the corresponding subalgebra of $\mathfrak{g}_1$).

Note that the group G_1 is simple according to Theorem 22. But since G_1 is simple, Q is a multiple of the Cartan-Killing form of $\mathfrak{g}_1$, therefore, Q is positive definite on $\mathfrak{g}_1$, and M is G_1 -normal. Theorem is proved. ■

We obviously get from Theorem 25 and Corollary 1

Corollary 10. *Every compact naturally reductive homogeneous Riemannian manifold with positive Euler characteristic is δ -homogeneous.*

According to Corollary 10, a compact naturally reductive homogeneous Riemannian manifold M , which is not δ -homogeneous, satisfies the condition $\chi(M) = 0$. In Section 15 we obtain examples of δ -homogeneous Riemannian manifolds with positive Euler characteristic, which are not normal homogeneous (consequently, are not naturally reductive).

9. ON ALGEBRAIC COROLLARIES OF THE δ -HOMOGENEITY

Let $(G/H, \mu)$ be a compact G - δ -homogeneous Riemannian manifold with a connected Lie group G , and let $\langle \cdot, \cdot \rangle$ be an $\text{Ad}(G)$ -invariant inner product on the Lie algebra $\mathfrak{g}$ of the group G . Denote by $\mathfrak{h}$ the Lie algebra of the group H , and consider some $\text{Ad}(H)$ -invariant complement $\mathfrak{p}$ to $\mathfrak{h}$ in $\mathfrak{g}$ (e.g., we can take $\mathfrak{p}$ from the $\langle \cdot, \cdot \rangle$ -orthogonal decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$). It is well known that the metric μ is generated by some $\text{Ad}(H)$ -invariant inner product $(\cdot, \cdot)$ on $\mathfrak{p}$, and there is the equality

$$(\cdot, \cdot) = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2} + \dots + x_s \langle \cdot, \cdot \rangle|_{\mathfrak{p}_s} \quad (9.6)$$

for some $\text{Ad}(H)$ -invariant pairwise orthogonal (with respect to both inner products) submodules $\mathfrak{p}_i$ ($1 \leq i \leq s$) of the $\text{Ad}(H)$ -module $\mathfrak{p}$ and for some positive numbers x_i ($1 \leq i \leq s$) such that $x_1 < x_2 < \dots < x_s$. Note that the modules $\mathfrak{p}_i$ need not to be $\text{Ad}(H)$ -irreducible.

For a vector $Z \in \mathfrak{g}$ let us denote by $Z_{\mathfrak{p}}$ and $Z_{\mathfrak{h}}$ its projections to subspaces $\mathfrak{p}$ and $\mathfrak{h}$ respectively, and for a vector $U \in \mathfrak{p}$ we will denote by U_i its projection to $\mathfrak{p}_i$, $1 \leq i \leq s$. The symbol $|\cdot|$ denotes the norm on $\mathfrak{p}$, generated by the scalar product $\langle \cdot, \cdot \rangle$.

We will give at first another simple proof of the fact that every (G) -normal homogeneous Riemannian manifold $(G/H, \mu)$ is (G) - δ -homogeneous. Let us consider for this the decomposition (9.6), where $s = 1$ and $x_1 = 1$. Choose any $X \in \mathfrak{p}$ and show that the vector X is δ -vector, see Definition 8. Let $a \in G$, then by $\text{Ad}(G)$ -invariance of the scalar product $\langle \cdot, \cdot \rangle$ we get

$$\langle \text{Ad}(a)(X), \text{Ad}(a)(X) \rangle = \langle X, X \rangle,$$

thus

$$|\text{Ad}(a)(X)|_{\mathfrak{p}}^2 = \langle \text{Ad}(a)(X)_{\mathfrak{p}}, \text{Ad}(a)(X)_{\mathfrak{p}} \rangle \leq \langle \text{Ad}(a)(X), \text{Ad}(a)(X) \rangle = \langle X, X \rangle = |X|^2.$$

Proposition 15 implies that $(G/H, \mu)$ is (G) - δ -homogeneous.

Now we derive some corollaries from δ -homogeneity of Riemannian manifolds in terms of Lie algebras.

Let us consider in a G - δ -homogeneous Riemannian manifold $(G/H, \mu)$ (with a closed connected transitive isometry group G) a geodesic γ , passing through the point eH in the direction V , $V \in \mathfrak{p} - \{0\}$. Suppose, that the Killing field $V + U$, $U \in \mathfrak{h}$ admits the maximum of its length on γ , and that this field generates an one-parameter motion group, one of whose orbit is γ (Theorem 10).

Proposition 20. *In the above condition the function $\varphi : G \rightarrow \mathbb{R}$, defined by the formula $\varphi(g) = |(\text{Ad}(g)(V + U))_{\mathfrak{p}}|$, where $g \in G$, has the absolute maximum at the point $g = e$.*

Corollary 11. *In the above condition one has the following:*

$$(V, [X, V + U]_{\mathfrak{p}}) = 0 \text{ for all } X \in \mathfrak{g}, \quad (9.7)$$

$$(V, [X, [X, V + U]]_{\mathfrak{p}}) + |[X, V + U]_{\mathfrak{p}}|^2 \leq 0 \text{ for all } X \in \mathfrak{g}. \quad (9.8)$$

Proof. Let us consider arbitrary $X \in \mathfrak{g}$. Then the function $f(t) = |(\text{Ad}(e^{tX})(V + U))_{\mathfrak{p}}|^2$ has its absolute maximum at the point $t = 0$. Now the statement of Corollary follows from the following:

$$f(t) = |V|^2 + 2(V, [X, V + U]_{\mathfrak{p}})t + ([X, V + U]_{\mathfrak{p}}|^2 + (V, [X, [X, V + U]]_{\mathfrak{p}}))t^2 + o(t^2) \text{ when } t \rightarrow 0.$$

■

Remark 11. Note that for $X \in \mathfrak{h}$ the relations (9.7) and (9.8) are fulfilled for any invariant metric.

Remark 12. The equation $(V, [X, V + U]_{\mathfrak{p}}) = 0$ in the previous corollary is a well known criterion for geodesic vectors (see Proposition 6).

Now we easily obtain

Theorem 26. *Let $(G/H, \mu)$ be a G - δ -homogeneous Riemannian manifold with connected Lie group G . Then for every $V \in \mathfrak{p}$ there is $U \in \mathfrak{h}$ such that for every $X \in \mathfrak{g}$ the following conditions fulfilled:*

$$(V, [X, V + U]_{\mathfrak{p}}) = 0, \quad (V, [X, [X, V + U]]_{\mathfrak{p}}) + |[X, V + U]_{\mathfrak{p}}|^2 \leq 0.$$

10. ON δ -HOMOGENEOUS MANIFOLD OF ONE SPECIAL TYPE

Let G be a compact connected Lie group, $H \subset K \subset G$ are its closed subgroup. Fix some $\text{Ad}(G)$ -invariant inner product $\langle \cdot, \cdot \rangle$ on the Lie algebra $\mathfrak{g}$ of the group G . Consider $\langle \cdot, \cdot \rangle$ -orthogonal decomposition

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p} = \mathfrak{h} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2,$$

where $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{p}_2$ is a Lie algebra of the group K . Obviously, $[\mathfrak{p}_2, \mathfrak{p}_1] \subset \mathfrak{p}_1$. Let μ be a G -invariant Riemannian metric on G/H , generated by the inner product

$$\langle \cdot, \cdot \rangle = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2}$$

on $\mathfrak{p}$ for some $x_1 > 0, x_2 > 0$ with $x_1 \neq x_2$.

For any vector $V \in \mathfrak{g}$ we denote by $V_{\mathfrak{p}}$ and $V_{\mathfrak{h}}$ its $(\langle \cdot, \cdot \rangle)$ -orthogonal projection to $\mathfrak{h}$ and $\mathfrak{p}$ respectively.

Proposition 21 ([43]). *Let $W = X + Y + Z$ be a geodesic vector on $(G/H, \mu)$, where $X \in \mathfrak{p}_1$, $Y \in \mathfrak{p}_2$, $Z \in \mathfrak{h}$. Then we have the following equalities:*

$$[Z, Y] = 0, \quad [X, Y] = \frac{x_1}{x_2 - x_1} [X, Z]. \quad (10.9)$$

Proof. By Theorem 26, for any $U \in \mathfrak{g}$ the equality $(X + Y, [U, X + Y + Z]_{\mathfrak{p}}) = 0$ holds. Therefore, we have

$$\begin{aligned} (X + Y, [U, X + Y + Z]_{\mathfrak{p}}) &= x_1 \langle X, [U, X + Y + Z] \rangle + x_2 \langle Y, [U, X + Y + Z] \rangle = \\ &= x_1 \langle [X + Y + Z, X], U \rangle + x_2 \langle [X + Y + Z, Y], U \rangle = \\ &= \langle (x_2 - x_1) [X, Y] + x_1 [Z, X] + x_2 [Z, Y], U \rangle = 0 \end{aligned}$$

for any $U \in \mathfrak{g}$. Since $[Z, Y] \in \mathfrak{p}_2$ and $[X, Y], [Z, X] \in \mathfrak{p}_1$, this proves Proposition. ■

Proposition 22. *Let $W = X + Y + Z$ be a δ -vector on $(G/H, \mu)$, where $X \in \mathfrak{p}_1$, $Y \in \mathfrak{p}_2$, $Z \in \mathfrak{h}$. Then for any $U \in \mathfrak{p}_1$ the following inequality holds:*

$$\begin{aligned} -x_1 \langle [U, X]_{\mathfrak{h}}, [U, X]_{\mathfrak{h}} \rangle + (x_2 - x_1) \langle [U, X]_{\mathfrak{p}_2}, [U, X]_{\mathfrak{p}_2} \rangle + (x_1 - x_2) \langle [U, Y], [U, X] \rangle + \\ (x_1 - x_2) \langle [U, Y], [U, Y] \rangle + x_1 \langle [U, X], [U, Z] \rangle + \\ (2x_1 - x_2) \langle [U, Y], [U, Z] \rangle + x_1 \langle [U, Z], [U, Z] \rangle \leq 0. \end{aligned} \quad (10.10)$$

Proof. According to Theorem 26 we get the inequality

$$(X + Y, [U, [U, X + Y + Z]_{\mathfrak{p}}]) + ([U, X + Y + Z]_{\mathfrak{p}}, [U, X + Y + Z]_{\mathfrak{p}}) \leq 0.$$

It is clear that $[Z, X], [Z, U], [Y, X], [Y, U] \in \mathfrak{p}_1$, $[Z, Y] \in \mathfrak{p}_2$. Therefore, using $\text{Ad}(G)$ -invariance of $\langle \cdot, \cdot \rangle$, we obtain

$$\begin{aligned} 0 &\geq (X + Y, [U, [U, X + Y + Z]_{\mathfrak{p}}]) + ([U, X + Y + Z]_{\mathfrak{p}}, [U, X + Y + Z]_{\mathfrak{p}}) = \\ &= -x_1 \langle [U, X], [U, X + Y + Z] \rangle - x_2 \langle [U, Y], [U, X + Y + Z] \rangle + \\ &= x_1 \langle [U, X]_{\mathfrak{p}_1} + [U, Y + Z], [U, X]_{\mathfrak{p}_1} + [U, Y + Z] \rangle + x_2 \langle [U, X]_{\mathfrak{p}_2}, [U, X]_{\mathfrak{p}_2} \rangle = \\ &= -x_1 \langle [U, X], [U, X] \rangle - x_1 \langle [U, X], [U, Y] \rangle - x_1 \langle [U, X], [U, Z] \rangle - x_2 \langle [U, Y], [U, X] \rangle - x_2 \langle [U, Y], [U, Y] \rangle \\ &= -x_2 \langle [U, Y], [U, Z] \rangle + x_1 \langle [U, X]_{\mathfrak{p}_1}, [U, X]_{\mathfrak{p}_1} \rangle + x_1 \langle [U, Y], [U, Y] \rangle + x_1 \langle [U, Z], [U, Z] \rangle \\ &= +2x_1 \langle [U, Y], [U, X] \rangle + 2x_1 \langle [U, X], [U, Z] \rangle + 2x_1 \langle [U, Y], [U, Z] \rangle + x_2 \langle [U, X]_{\mathfrak{p}_2}, [U, X]_{\mathfrak{p}_2} \rangle = \\ &= -x_1 \langle [U, X]_{\mathfrak{h}}, [U, X]_{\mathfrak{h}} \rangle + (x_2 - x_1) \langle [U, X]_{\mathfrak{p}_2}, [U, X]_{\mathfrak{p}_2} \rangle + (x_1 - x_2) \langle [U, Y], [U, X] \rangle + \\ &= (x_1 - x_2) \langle [U, Y], [U, Y] \rangle + x_1 \langle [U, X], [U, Z] \rangle + (2x_1 - x_2) \langle [U, Y], [U, Z] \rangle + x_1 \langle [U, Z], [U, Z] \rangle, \end{aligned}$$

which proves Proposition. ■

Corollary 12. *If in conditions of Proposition 22 $X = 0$, then for any $U \in \mathfrak{p}_1$ we have*

$$(x_1 - x_2) \langle [U, Y], [U, Y] \rangle + (2x_1 - x_2) \langle [U, Y], [U, Z] \rangle + x_1 \langle [U, Z], [U, Z] \rangle \leq 0. \quad (10.11)$$

Proposition 23. *For any δ -vector $X + Y + Z$ on $(G/H, \mu)$ the vector $Y + Z$ is a δ -vector on K/H (with the induced metric). In particular, if $(G/H, \mu)$ is G - δ -homogeneous, then K/H with the induced metric is K - δ -homogeneous.*

Proof. For any $\text{Ad}(a)$, where $a \in K$, we have $\text{Ad}(a)(\mathfrak{p}_1) = \mathfrak{p}_1$. Moreover, $\text{Ad}(a)|_{\mathfrak{p}_1}$ is orthogonal transformation. Since

$$(X, X) + (Y, Y) = (X + Y, X + Y) \geq (\text{Ad}(a)(X + Y + Z)|_{\mathfrak{p}}, \text{Ad}(a)(X + Y + Z)|_{\mathfrak{p}}) = \\ (X, X) + (\text{Ad}(a)(Y + Z)|_{\mathfrak{p}}, \text{Ad}(a)(Y + Z)|_{\mathfrak{p}})$$

for any $a \in K$, the vector $Y + Z$ is δ -vector for K/H . Remark that really the Riemannian subspace K/H of $(G/H, \mu)$ is K -normal, because $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{p}_2$. ■

Proposition 24. *For any geodesic vector $X + Y + Z$ on $(G/H, \mu)$ the vector $Y + Z$ is geodesic vector on K/H (with the induced metric).*

Proof. By Proposition 6, $X + Y + Z$ is geodesic if and only if for any $U \in \mathfrak{g}$ we have $(X + Y, [U, X + Y + Z]_{\mathfrak{p}}) = 0$. Let $U \in \mathfrak{p}_2 \oplus \mathfrak{h}$, then $[U, X + Y + Z]_{\mathfrak{p}_1} = [U, X]$, $[U, X + Y + Z]_{\mathfrak{p}_2} = [U, Y + Z]_{\mathfrak{p}}$. Therefore, we have $(Y, [U, Y + Z]_{\mathfrak{p}}) = 0$, since $(X, [U, X]) = 0$. Since $U \in \mathfrak{h} \oplus \mathfrak{p}_2$ may be arbitrary, we get that the vector $Y + Z$ is a geodesic vector on K/H . ■

Proposition 25. *If vectors $\tilde{X} + Y + Z$ and $X + Y + Z$ both are δ -vectors on $(G/H, \mu)$, then*

$$x_1 \langle [\tilde{X}, X]_{\mathfrak{h}}, [\tilde{X}, X]_{\mathfrak{h}} \rangle \geq (x_2 - x_1) \langle [\tilde{X}, X]_{\mathfrak{p}_2}, [\tilde{X}, X]_{\mathfrak{p}_2} \rangle.$$

Proof. From Proposition 21 we have the equality $[\tilde{X}, Y] = x_1/(x_2 - x_1)[\tilde{X}, Z]$. Putting $U = \tilde{X}$ in the inequality (10.10) and using the above equality, we prove Proposition. ■

Proposition 26. *Suppose that $(G/H, \mu)$ is G - δ -homogeneous. Let $X \in \mathfrak{p}_1$, $Y \in \mathfrak{p}_2$, $a = \exp(tY)$ for some $t \in \mathbb{R}$, $\tilde{X} = \text{Ad}(a)(X)$. Then the following inequality holds:*

$$x_1 \langle [\tilde{X}, X]_{\mathfrak{h}}, [\tilde{X}, X]_{\mathfrak{h}} \rangle \geq (x_2 - x_1) \langle [\tilde{X}, X]_{\mathfrak{p}_2}, [\tilde{X}, X]_{\mathfrak{p}_2} \rangle.$$

Proof. Let $Z \in \mathfrak{h}$ be such a vector that $X + Y + Z$ is δ -vector. From Proposition 21 we have $[Z, Y] = 0$. This implies that $\text{Ad}(a)(Z) = Z$. Besides this, $\text{Ad}(a)(Y) = Y$, and $(X, X) = (\tilde{X}, \tilde{X})$, since $\text{Ad}(a)|_{\mathfrak{p}_1}$ is $(\cdot, \cdot)$ -orthogonal. Therefore, the vector $\tilde{X} + Y + Z = \text{Ad}(a)(X + Y + Z)$ is δ -vector too. Now we can apply Proposition 25. ■

Since for $a = \exp(tY)$ we have $\text{Ad}(a)(X) = X + [Y, X]t + o(t)$ when $t \rightarrow 0$, we get the following infinitesimal version of Proposition 26.

Proposition 27. *Suppose that $(G/H, \mu)$ is G - δ -homogeneous. Let $X \in \mathfrak{p}_1$, $Y \in \mathfrak{p}_2$, then the following inequality holds:*

$$x_1 \langle [[Y, X], X]_{\mathfrak{h}}, [[Y, X], X]_{\mathfrak{h}} \rangle \geq (x_2 - x_1) \langle [[Y, X], X]_{\mathfrak{p}_2}, [[Y, X], X]_{\mathfrak{p}_2} \rangle.$$

11. ROOT SYSTEMS OF COMPACT SIMPLE LIE ALGEBRAS

We give here some information about root systems of a compact simple Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle = -B)$ with the Killing form B , which can be find in books [23, 14].

The Lie algebra $\mathfrak{g}$ admits a direct $\langle \cdot, \cdot \rangle$ -orthogonal decomposition $\mathfrak{t} \oplus \text{Lin}\{\cup_{\alpha \in \Delta} V_{\alpha}\}$ into (non-zero) vector subspaces, where $\alpha \in \mathfrak{t}^*$ is some (non-zero) real-valued linear form on the Cartan subalgebra $\mathfrak{t}$ of Lie algebra $\mathfrak{g}$, $V_{\alpha} = V_{-\alpha}$ is some 2-dimensional $\text{ad}(\mathfrak{t})$ -invariant vector subspace, and Lin means a linear span. Using the restriction (of non-degenerate) inner product $\langle \cdot, \cdot \rangle$ to $\mathfrak{t}$, we will naturally identify α with vector in $\mathfrak{t}$. All such forms (vectors) α are called *roots* of Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$, and the set Δ of all such roots α is called *root system* of Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$. It is easy to see that $[V_{\alpha}, V_{\alpha}]$ is one-dimensional subalgebra of $\mathfrak{t}$ spanned on the root α , and $[V_{\alpha}, V_{\alpha}] \oplus V_{\alpha}$ is a Lie algebra isomorphic to $\mathfrak{su}(2)$. This implies that vector subspaces $V_{\alpha}, \alpha \in \Delta$, admit bases $\{u_{\alpha}, v_{\alpha}\}$ with the following commutator relations

$$[h, u_{\alpha}] = -\langle \alpha, h \rangle v_{\alpha}, \quad [h, v_{\alpha}] = \langle \alpha, h \rangle u_{\alpha}, \quad h \in \mathfrak{t}, \quad [u_{\alpha}, v_{\alpha}] = -\frac{4}{\langle \alpha, \alpha \rangle} \alpha. \quad (11.12)$$

Moreover, for $\alpha \neq \pm\beta$,

$$[u_{\alpha}, u_{\beta}] = N_{\alpha, \beta} u_{\alpha + \beta} + N_{\alpha, -\beta} u_{\alpha - \beta}, \quad (11.13)$$

$$[v_\alpha, v_\beta] = -N_{\alpha,\beta}u_{\alpha+\beta} + N_{\alpha,-\beta}u_{\alpha-\beta}, \quad (11.14)$$

$$[u_\alpha, v_\beta] = N_{\alpha,\beta}v_{\alpha+\beta} - N_{\alpha,-\beta}v_{\alpha-\beta}, \quad (11.15)$$

$$[v_\alpha, u_\beta] = N_{\alpha,\beta}v_{\alpha+\beta} + N_{\alpha,-\beta}v_{\alpha-\beta}, \quad (11.16)$$

where (integer) numbers $N_{\alpha,\beta}$ are defined as follows:

$$N_{-\alpha,-\beta} = N_{\alpha,\beta}, \quad N_{\alpha,\beta} = \pm(q+1)$$

for $\alpha, \beta, \alpha + \beta \in \Delta$, where q is the greatest integer number j such that $\beta - j\alpha \in \Delta$. We suggest in these formulas that $N_{\gamma,\delta} = 0$, if $\gamma + \delta$ is not a root. From (11.12) and the invariance of $\langle \cdot, \cdot \rangle$ with respect to automorphisms of $\mathfrak{g}$, it is easy to obtain

$$\langle u_\alpha, u_\alpha \rangle = \langle v_\alpha, v_\alpha \rangle = \frac{4}{\langle \alpha, \alpha \rangle}. \quad (11.17)$$

The formulas above imply

Lemma 4.

$$[V_\alpha, V_\beta] = V_{\alpha+\beta} + V_{\alpha-\beta}.$$

The root system Δ is invariant relative to the Weyl group $W = W(T)$. Besides this:

(i) For every root $\alpha \in \Delta \subset \mathfrak{t}$ the Weyl group W contains the reflection φ_α in the plane P_α , which is orthogonal to the root α (with respect to $\langle \cdot, \cdot \rangle$).

(ii) Reflections from (i) generate W .

We list below the root systems of that simple compact Lie groups which we shall need later:

$$\begin{aligned} A_l &: e_i - e_j, \quad i \neq j, \quad i, j = 0, 1, \dots, l. \\ B_l &: \pm e_i, \quad i = 1, 2, \dots, l; \quad \pm e_i \pm e_j, \quad i < j, \quad i, j = 1, 2, \dots, l. \\ C_l &: \pm 2e_i, i = 1, 2, \dots, l; \quad \pm e_i \pm e_j, \quad i < j, \quad i, j = 1, 2, \dots, l. \\ D_l &: \pm e_i \pm e_j, \quad i < j, \quad i, j = 1, 2, \dots, l. \\ g_2 &: e_i - e_j; \quad \pm \left(\sum_{i=1}^3 e_i - 3e_j \right), \quad i, j = 1, 2, 3. \\ f_4 &: \pm e_i, \quad \pm e_i \pm e_j, \quad \frac{1}{2}(\pm e_1 \pm e_2 \pm e_3 \pm e_4), \quad i, j = 1, 2, 3, 4. \end{aligned}$$

Here $A_{l-1} = su(l)$, $B_l = so(2l+1)$, $C_l = sp(l)$, $D_l = so(2l)$. Let us remark that all roots of any Lie algebra A_l, D_l, e_6, e_7, e_8 have one and the same lengths. The roots of any other simple Lie algebra have two different lengths, so we have the systems $\Delta_l \subset \Delta$ and $\Delta_s \subset \Delta$ of all long and short roots respectively. If $\alpha \in \Delta_l, \beta \in \Delta_s$ for B_l, C_l, f_4 (respectively g_2), then $|\alpha| = \sqrt{2}|\beta|$ (respectively $|\alpha| = \sqrt{3}|\beta|$), where $|X| = \sqrt{\langle X, X \rangle}$. In all cases two roots of equal length may constitute the angles $\frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}$. The roots of different length for B_l, C_l, f_4 (respectively, g_2) may constitute the angles $\frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}$ (respectively $\frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$).

By theorem 20, all simply connected homogeneous spaces G/H of positive Euler characteristic with a simple Lie group G are in one-to one correspondence with Lie subalgebras $\mathfrak{h}$, such that $\mathfrak{t} \subset \mathfrak{h} \subset \mathfrak{g}$ and $\mathfrak{h} \neq \mathfrak{g}$; we must identify subalgebras, which are $\text{Ad}(g)$ -conjugate with respect to some $g \in G$ such that $\text{Ad}(g)(\mathfrak{t}) = \mathfrak{t}$. Any such Lie subalgebra $\mathfrak{h}$ is defined by a class of pairwise W -isomorphic closed symmetric root subsystems A of Δ , not equal to Δ . By definition, $A \subset \Delta$ is *closed*, if $\alpha, \beta \in A$ and $\alpha \pm \beta \in \Delta$ imply $\alpha \pm \beta \in A$, and *symmetric*, if $-\alpha \in A$ together with $\alpha \in A$. Then

$$\mathfrak{h} = \mathfrak{t} \oplus \text{Lin}\{\cup_{\alpha \in A} V_\alpha\}, \quad (11.18)$$

where Lin means a linear span.

12. ON THE GROUP G_2

Let's describe all simply connected homogeneous spaces G/H of positive Euler characteristic for $G = G_2 = \text{Aut}(\mathbb{C}a)$. For this we use the considerations from the previous section.

Let us give a description of the root system Δ of the Lie algebra g_2 . There are two simple roots $\alpha, \beta \in \Delta$ such that $\angle(\alpha, \beta) = \frac{5\pi}{6}$ and $|\alpha| = \sqrt{3}|\beta|$. Then

$$\Delta = \{\pm\alpha, \pm\beta, \pm(\alpha + \beta), \pm(\alpha + 2\beta), \pm(\alpha + 3\beta), \pm(2\alpha + 3\beta)\}.$$

Under this, $\pm\alpha, \pm(\alpha + 3\beta), \pm(2\alpha + 3\beta)$ are all long roots. One can easily see that all non W -isomorphic closed symmetric root subsystems of Δ_{G_2} , not equal to Δ_{G_2} , are $\emptyset, \{\pm\alpha\}, \{\pm\beta\}, \{\pm\beta, \pm(2\alpha + 3\beta)\}, \{\pm\alpha, \pm(\alpha + 3\beta), \pm(2\alpha + 3\beta)\}$.

The first three cases give us respectively the following (generalized) flag manifolds: G_2/T^2 , $G_2/SU(2)SO(2)$, and $G_2/A_{1,3}SO(2)$, where $A_{1,3}$ is a Lie group with Lie subalgebra of the type A_1 of index 3, see [37]. D.V. Alekseevsky and A. Arvanitoyeorgos proved in [4] that all G_2 -invariant Riemannian g.o. metrics on them with the full connected isometry group G_2 are G_2 -normal. The discussion in Section 8 implies that any G_2 -invariant metric on these spaces, whose full connected isometry group is not G_2 , is $SO(7)$ -normal (symmetric) metric on $G_2/SU(2) \cdot SO(2) = Gr_{7,2}^+$.

The last two closed symmetric root subsystems are maximal, so they correspond to maximal Lie subalgebras in g_2 , which are respectively isomorphic to $su(2) \oplus su(2)$ and $su(3)$ with the corresponding compact connected Lie subgroups $SO(4)$ and $SU(3)$ and homogeneous spaces $G_2/SO(4)$ and $G_2/SU(3) = S^6$, compare with [37]. In the first (second) case

$$\mathfrak{p} = V_\alpha \oplus V_{\alpha+\beta} \oplus V_{\alpha+2\beta} \oplus V_{\alpha+3\beta}$$

(respectively

$$\mathfrak{p} = V_\beta \oplus V_{\alpha+\beta} \oplus V_{\alpha+2\beta}).$$

It's well-known that irreducible components of a representation of a compact Lie algebra are uniquely determined up to equivalence. As a corollary, applying this to the adjoint representation of Lie subalgebra $\mathfrak{t} \subset \mathfrak{h}$ on $\mathfrak{p}$, one get that for any $\text{ad}(\mathfrak{h})$ -invariant subspace $V \subset \mathfrak{p}$ there exists an equivalent $\text{ad}(\mathfrak{h})$ -invariant subspace $V' \subset \mathfrak{p}$, which is a direct sum of the given root vector subspaces $V_\gamma, \gamma \in R$. One can easily see that in both cases above there is no such $\text{ad}(\mathfrak{h})$ -invariant subspace $V' \subset \mathfrak{p}$ besides $\mathfrak{p}$ and $\{0\}$. Thus the space $\mathfrak{p}$ is $\text{ad}(\mathfrak{h})$ -irreducible. This means that the corresponding homogeneous spaces G_2/H are strongly isotropy irreducible. Then any G_2 -invariant Riemannian metric on G_2/H is G_2 -normal.

Remark 13. Note that $G_2/SO(4)$ is irreducible symmetric space, see [12].

Therefore, we have the following

Proposition 28. *Any g.o. (any δ -homogeneous, in particular) Riemannian homogeneous manifold $(G_2/H, \mu)$ of positive Euler characteristic is either G_2 -normal or $SO(7)$ -normal.*

Let us remark at the end that the very last root subsystem contains only the long roots.

13. CALCULATIONS WITH ROOTS

Let suppose that in the Notation of Section 6, $(M = G/H, \mu)$ is G - δ -homogeneous simply connected indecomposable Riemannian manifold with positive Euler characteristic. Then G is simple by Theorem 22, and we have inclusions $T \subset H \subset G$, where T is a maximal torus in G . Then we have some $\text{Ad}(T)$ -invariant $\langle \cdot, \cdot \rangle$ -orthogonal decomposition

$$\mathfrak{g} = \mathfrak{t} \oplus \text{Lin}\{\cup_{\gamma \in C} V_\gamma\} \oplus \text{Lin}\{\cup_{\alpha \in D} V_\alpha\},$$

$C \cup D = \Delta$ is a set of all roots for Lie group G with respect to Lie algebra $\mathfrak{t}$ of T , $V_\alpha = V_{-\alpha}$ and $V_\gamma = V_{-\gamma}$ are two-dimensional "root spaces", and the first two summands give us a decomposition of the Lie algebra $\mathfrak{h}$ of the Lie group H , the last summand gives $\text{Ad}(H)$ -invariant vector subspace $\mathfrak{p}$.

Proposition 29. *Let $\alpha_1, \dots, \alpha_k \in D$ are linearly independent roots. Then there is a unique (up to multiplication by constant) vector $t_c \in \text{Lin}\{\alpha_1, \dots, \alpha_k\}$ such that for some real number s , $\text{Ad}(\exp(st_c)) = -\text{Id}$ on $\oplus_{i=1}^k V_{\alpha_i}$.*

Proof. One can easily prove this by using the dual basis in Euclidean space $\text{Lin}\{\alpha_1, \dots, \alpha_k\}$. ■

Proposition 30. *Let $\alpha_1, \dots, \alpha_k \in D$ are linearly independent roots and $v = \sum_{i=1}^k v_i$, where $v_i \in V_{\alpha_i}$, $i = 1, \dots, k$, are non-zero vectors. Let $u(v) \neq 0$, (see Remark 7) and C_v is the set of all $\gamma \in C$ such that V_γ -component of $u(v)$ is not zero. Then*

$$C_v \neq \emptyset, \quad C_v \subset \text{Lin}\{\alpha_1, \dots, \alpha_k\} - t_c^\perp,$$

where $t_c^\perp$ is the orthogonal complement in $\text{Lin}\{\alpha_1, \dots, \alpha_k\}$ to the vector t_c from Proposition 29.

Proof. Really, if $C_v = \emptyset$, then $u(v) := u \in t$ and by Proposition 29

$$\text{Ad}(\exp(st_c))(w) = -w, \quad \text{Ad}(\exp(st_c))(u(w)) = u(w), \quad (13.19)$$

since $[u, t_c] = 0$. This contradicts to Proposition 14. So, $C_v \neq \emptyset$.

Now, if some $\gamma \in C_v$ is not in $\text{Lin}\{\alpha_1, \dots, \alpha_k\}$, then one can find a vector $w \in t$, which is orthogonal to all $\alpha_1, \dots, \alpha_k$, but $\langle w, \gamma \rangle \neq 0$. Then $[w, v] = 0$, while $[w, u(v)] \neq 0$ which contradicts to Proposition 14.

Finally, if $C_v \in t_c^\perp$, then one more we have (13.19), which is impossible by Proposition 14. ■

Since roots $\alpha \in D$, $\gamma \in C$ are non-collinear, the next proposition follows from Propositions 14 and 30.

Proposition 31. *If $v \in V_\alpha$, $\alpha \in D$, then $\|v\| = |v| = \sqrt{(v, v)}$, i.e. v is a δ -vector.*

Proposition 32. *We have at most two possibilities: $(\cdot, \cdot) = x\langle \cdot, \cdot \rangle$ on $\mathfrak{p}$ or we have an $\text{Ad}(H)$ -invariant $\langle \cdot, \cdot \rangle$ -orthogonal direct decomposition $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$ such that $(\cdot, \cdot) = x_1\langle \cdot, \cdot \rangle$ on $\mathfrak{p}_1$ and $(\cdot, \cdot) = x_2\langle \cdot, \cdot \rangle$ on $\mathfrak{p}_2$, where $x_1 \neq x_2$. We have necessarily the first possibility, if all roots of G have one and the same length.*

Proof. The elements $\text{Ad}(n)$, $n \in N(T)$, generate on $\mathfrak{t}$ a finite Weyl group $W = W(T)$. It is known that W is generated by orthogonal reflections in hyper-planes in $\mathfrak{t}$, orthogonal to roots in $\Delta \subset \mathfrak{t}$. From this and known classifications of roots systems of compact simple Lie groups one can easily deduce that W acts transitively on every set of roots of equal lengths. There are at most two such sets in Δ : the set of all short roots Δ_s and the set of all long roots Δ_l (see Section 11). At the same time $\text{Ad}(n)$, $n \in N(T)$, acts transitively on the set of root vector spaces V_α , $\alpha \in \Delta_l$ or $\alpha \in \Delta_s$. Since $\|\cdot\|$ and $\langle \cdot, \cdot \rangle$ are $\text{Ad}(G)$ -invariant, we get by Proposition 31 that

$$(v_\alpha, v_\alpha) = \|v_\alpha\|^2 = \|v_\beta\|^2 = (v_\beta, v_\beta)$$

and

$$\langle v_\alpha, v_\alpha \rangle = \langle v_\beta, v_\beta \rangle,$$

if $\alpha, \beta \in \Delta_l$ or $\alpha, \beta \in \Delta_s$. Here $v_\alpha \in V_\alpha$ mean special vectors from Section 11. From this follow the required statements. ■

Corollary 13. *Any G - δ -homogeneous Riemannian manifold $(G/H, \mu)$ of positive Euler characteristic with $G = SU(l+1)$, $SO(2l)$, E_6 , E_7 , or E_8 is G -normal.*

Therefore, we should examine only the second case in Proposition 32. Later on we shall use the following notation in this case:

$$\mathfrak{p}_1 = \text{Lin}\{\cup_{\beta \in B} V_\beta\}, \quad \mathfrak{p}_2 = \text{Lin}\{\cup_{\alpha \in A} V_\alpha\}, \quad A = \Delta_l \cap D, \quad B = \Delta_s \cap D, \quad (13.20)$$

where Δ_l (Δ_s) means the set of long (respectively, short) roots of Lie algebra $\mathfrak{g}$.

Lemma 5. *Let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2$ as above, then $[\mathfrak{p}_1, \mathfrak{p}_2] \neq 0$.*

Proof. Let us suppose that $[\mathfrak{p}_1, \mathfrak{p}_2] = 0$ and show that in this case $\mathfrak{q} := \mathfrak{p}_1 + [\mathfrak{p}_1, \mathfrak{p}_1]$ is an proper ideal of $\mathfrak{g}$. For this goal it is sufficient to show that $[\mathfrak{h}, \mathfrak{q}] \subset \mathfrak{q}$, $[\mathfrak{p}_1, \mathfrak{q}] \subset \mathfrak{q}$ and $[\mathfrak{p}_2, \mathfrak{q}] \subset \mathfrak{q}$.

Since $[\mathfrak{h}, \mathfrak{p}_1] \subset \mathfrak{p}_1$ and $[\mathfrak{p}_2, \mathfrak{p}_1] = 0$, then by the Jacobi identity we get $[h, [\mathfrak{p}_1, \mathfrak{p}_1]] \subset [[h, \mathfrak{p}_1], \mathfrak{p}_1] \subset [\mathfrak{p}_1, \mathfrak{p}_1] \subset \mathfrak{q}$ and $[\mathfrak{p}_2, [\mathfrak{p}_1, \mathfrak{p}_1]] = 0$. Therefore, $[\mathfrak{h}, \mathfrak{q}] \subset \mathfrak{q}$ and $[\mathfrak{p}_2, \mathfrak{q}] = 0$.

For any $X, Y \in \mathfrak{p}_1$ and $Z \in \mathfrak{p}_2$ we have $\langle [X, Y], Z \rangle = -\langle Y, [X, Z] \rangle = 0$, since $[\mathfrak{p}_1, \mathfrak{p}_2] = 0$. Hence, $[\mathfrak{p}_1, \mathfrak{p}_1] \subset \mathfrak{p}_1 \oplus \mathfrak{h}$, and $[\mathfrak{p}_1, \mathfrak{q}] \subset [\mathfrak{p}_1, \mathfrak{p}_1] + [\mathfrak{p}_1, [\mathfrak{p}_1, \mathfrak{p}_1]] \subset [\mathfrak{p}_1, \mathfrak{p}_1] + [\mathfrak{p}_1, \mathfrak{h}] \subset \mathfrak{q}$.

Consequently, $\mathfrak{q}$ is an ideal of $\mathfrak{g}$. This ideal is proper, since $\mathfrak{p}_2$ is $\langle \cdot, \cdot \rangle$ -ortogonal to $\mathfrak{q}$ (see above). On the other hand, $\mathfrak{g}$ is a simple Lie algebra and contains no nontrivial ideal. This contradiction proves Lemma. ■

Lemma 6. *Let suppose that the root system Δ of a compact simple Lie algebra $\mathfrak{g} \neq g_2$ contains two roots $\alpha \in \Delta_l$, $\beta \in \Delta_s$ of different lengths. Then at most one of $\alpha + \beta$ or $\alpha - \beta$ is a root in Δ .*

Proof. By previous description of Δ , we have exactly three possibilities for the angle between α and β : $\frac{\pi}{4}$, $\frac{\pi}{2}$, $\frac{3\pi}{4}$. In the second case no one of terms $\alpha + \beta$ or $\alpha - \beta$ is a root. Otherwise there would be a root, longer than α , which is impossible. In the first (respectively, third) case $\alpha - \beta$ (respectively, $\alpha + \beta$) is a root, but not $\alpha + \beta$ (respectively $\alpha - \beta$). ■

Lemma 7. 1) *The vector subspace*

$$\eta = \mathfrak{t} \oplus \text{Lin}\{\cup_{\alpha \in \Delta_l} V_\alpha\}$$

is a Lie subalgebra in $\mathfrak{g}$.

2) *The vector subspace η is a maximal subalgebra in $\mathfrak{g}$, if $G \neq F_4$ and $G \neq Sp(l)$, $l \geq 3$.*

3) *If $G = Sp(l)$, then all non-collinear roots in Δ_l are mutually orthogonal and $[V_{\alpha_1}, V_{\alpha_2}] = 0$, if $\alpha_1 \neq \pm \alpha_2$ are roots in Δ_l .*

4) *If $G = F_4$, then η is isomorphic to $so(8) = spin(8)$. There is $\text{ad}(\eta)$ -invariant decomposition*

$$\mathfrak{g} = f_4 = \eta \oplus \mathfrak{q}_1 \oplus \mathfrak{q}_2 \oplus \mathfrak{q}_3,$$

where $\mathfrak{q}_3 = \text{Lin}\{\cup_{\beta \in \Delta_a} V_\beta\}$, and Δ_a consists of all roots in Δ_s of a form $\pm e_i$, $i = 1, 2, 3, 4$; $\mathfrak{q}_1$ ($\mathfrak{q}_2$) is spanned on the root spaces of roots of the form $1/2(\pm e_1 \pm e_2 \pm e_3 \pm e_4)$ (see Section 11) with the odd (respectively, even) number of signs “-” in this formula. All modules $\mathfrak{q}_i$ are $\text{ad}(\eta)$ -irreducible, and $\mathfrak{r}_i = \eta \oplus \mathfrak{q}_i$, $1 \leq i \leq 3$, is a Lie algebra isomorphic to $so(9) = spin(9)$. For $i \neq j$ there is an automorphism of f_4 preserving η and $\mathfrak{t}$, which maps $\mathfrak{r}_i$ to $\mathfrak{r}_j$. Any proper subalgebra of $\mathfrak{g} = f_4$, containing η and different from η , is one of the subalgebra $\mathfrak{r}_i$, $1 \leq i \leq 3$.

Proof. For $G = G_2$ all statements can be checked directly and easily.

Let G be another simple group (with roots of different lengths), and $\alpha, \beta \in \Delta_l$. Then $\langle \alpha, \beta \rangle = 0$ or $\angle(\alpha, \beta) = \frac{2\pi}{3}$ or $\angle(\alpha, \beta) = \frac{\pi}{3}$. In the first case $\alpha \pm \beta$ cannot be a roots, so $[V_\alpha, V_\beta] = 0$. In the second (third) case orthogonal reflection of $\mathfrak{t}$ in the hyperplane, $\langle \cdot, \cdot \rangle$ -orthogonal to α (respectively, $-\alpha$), maps the root β to the vector $\alpha + \beta$ (respectively, to $\beta - \alpha$), so this vector is a long root. At the same time, $\alpha - \beta$ (respectively, $\beta + \alpha$) is not a root. So, we get $[V_\alpha, V_\beta] = V_{\alpha+\beta}$ (respectively, $[V_\alpha, V_\beta] = V_{\alpha-\beta}$). This finished the proof of the first statement.

The second statement easily follows from the list of all roots of a simple Lie algebra.

Let us remark that any maximal subalgebra θ in $\mathfrak{g} = sp(l)$, $l \geq 3$, (with root system C_l), containing η , has a form

$$\theta = \eta \oplus \text{Lin}\{\cup_{\alpha \in \Delta_s - \Delta_i} V_\alpha\},$$

where Δ_i contains all roots in Δ_s of a form $\pm e_i \pm e_j$ for a fixed $1 \leq i \leq l$, and all $j \neq i$. All these Lie algebras θ_i are mutually isomorphic under automorphisms of $\mathfrak{g}$ and are isomorphic to the Lie algebra $\theta_1 = sp(1) \oplus sp(l-1)$. So, if Θ is compact connected Lie subgroup in

$G = Sp(l)$ with Lie algebra θ_1 , then we get the homogeneous space $G/\Theta = Sp(l)/Sp(1) \times Sp(l-1) = \mathbb{H}P^{(l-1)}$.

All long roots for Lie algebra $sp(l)$ has the form $\pm 2e_i, 1 \leq i \leq l$, so we get the third statement.

One can check the first three statements of 4) directly. All other statement are proved in [2]. ■

Lemma 8. *The module $\mathfrak{k} := \mathfrak{h} \oplus \mathfrak{p}_2$ (see (13.20)) is a Lie subalgebra of $\mathfrak{g}$. As a corollary, $[\mathfrak{p}_2, \mathfrak{p}_1] \subset \mathfrak{p}_1$.*

Proof. It is clear that $[\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{h} \subset \mathfrak{k}$, $[\mathfrak{h}, \mathfrak{p}_2] \subset \mathfrak{p}_2 \subset \mathfrak{k}$ and $\eta \subset \mathfrak{k}$ (see Lemma 7). Note also that $[\mathfrak{p}_2, \mathfrak{p}_2] \subset [\eta, \eta] \subset \eta \subset \mathfrak{k}$. These considerations prove the first statement. The second statement is evident. ■

The previous Lemma permits now to use all results of Section 10.

Proposition 33. *Let suppose that we have the second possibility in Proposition 32 (so Δ has roots of two different lengths), and $\mathfrak{g} \neq g_2$. There are $\alpha \in A$, $\beta \in B$ (see (13.20)) such that $[V_\alpha, V_\beta] \neq 0$. For any such α, β , either $\alpha + 2\beta \in C$ or $\alpha - 2\beta \in C$. Moreover, the following inequality holds:*

$$x_1 < x_2 \leq 2x_1. \quad (13.21)$$

Proof. The first statement follows from Lemma 5.

If $[V_\alpha, V_\beta] \neq 0$, then by Lemma 6 we have only two possible cases for the angle between α and β : $\frac{\pi}{4}$ or $\frac{3\pi}{4}$. Both cases are quite similar, so let us consider the second one. In this case

$$\alpha + \beta \in \Delta_s, \alpha + 2\beta \in \Delta_l, \quad |\alpha|_1 = |\alpha + 2\beta|_1 = \sqrt{2}|\beta|_1 = \sqrt{2}|\alpha + \beta|_1, \quad (13.22)$$

where $|X|_1 = \sqrt{\langle X, X \rangle}$ for $X \in \mathfrak{g}$. Then

$$[[u_\alpha, u_\beta], u_\beta] = [N_{\alpha, \beta} u_{\alpha + \beta}, u_\beta] = N_{\alpha, \beta} (N_{\alpha + \beta, -\beta} u_\alpha + N_{\alpha + \beta, \beta} u_{\alpha + 2\beta}).$$

Here $N_{\alpha, \beta} = \pm(q+1)$, where $q = \max\{j : \beta - j\alpha \in \Delta\} = 0$, so $N_{\alpha, \beta} = \pm 1$. $N_{\alpha + \beta, -\beta} = \pm(p+1)$, where $p = \max\{j : -\beta - j(\alpha + \beta) \in \Delta\} = 1$, so $N_{\alpha + \beta, -\beta} = \pm 2$. $N_{\alpha + \beta, \beta} = \pm(l+1)$, where $l = \max\{j : \beta - j(\alpha + \beta) \in \Delta\} = 1$, so $N_{\alpha + \beta, \beta} = \pm 2$. Hence we get

$$[[u_\alpha, u_\beta], u_\beta] = 2(\pm u_\alpha \pm u_{\alpha + 2\beta}), \quad (13.23)$$

where one needs to take only one of four possible choices of signs. Since $u_\alpha \in \mathfrak{p}_2$, we see from (13.23) that $[[u_\alpha, u_\beta], u_\beta]_{\mathfrak{p}_2} \neq 0$. Then Proposition 27 implies that $[[u_\alpha, u_\beta], u_\beta]_{\mathfrak{h}} \neq 0$. It follows from the formula (13.23) that $[[u_\alpha, u_\beta], u_\beta]_{\mathfrak{h}} = \pm 2u_{\alpha + 2\beta}$. Hence $\alpha + 2\beta \in C$.

In order to prove the inequality $x_1 < x_2$ take a δ -vector $Y \in \mathfrak{p}_2$ and some $U \in \mathfrak{p}_1$ such that $[U, Y] \neq 0$ (it is possible according to Proposition 31 and Lemma 5). Using the inequality (10.11) of Corollary 12 in this case, we get $(x_1 - x_2)\langle [U, Y], [U, Y] \rangle \leq 0$, therefore, $x_1 < x_2$.

It is possible to prove the inequality $x_2 \leq 2x_1$, using Proposition 27. But we give a more clear proof. Let's consider the (Ad(G)-invariant) Chebyshev's norm $\|\cdot\|$ on $\mathfrak{g}$, corresponding to G - δ -homogeneous space $(G/H, \mu)$ (see Theorem 6). According to Proposition 31, for any root $\alpha \in A$ every $X \in V_\alpha$ is a δ -vector. Therefore, $\|X\| = \sqrt{\langle X, X \rangle} = \sqrt{x_2}|X|_1$. Similarly, for any root $\beta \in B$ every $Y \in V_\beta$ is a δ -vector and $\|Y\| = \sqrt{\langle Y, Y \rangle} = \sqrt{x_1}|Y|_1$. By above argument we can suppose that (13.22) is satisfied. Using the equations (11.12), (13.22) and Ad(G)-invariance of $\|\cdot\|$ and $|\cdot|_1$, we get that $\|\alpha\| = \sqrt{x_2}|\alpha|_1 = \sqrt{2x_2}|\beta|_1$ and $\|\beta\| = \sqrt{x_1}|\beta|_1$. According to Ad(G)-invariance of $\|\cdot\|$ and $|\cdot|_1$ we get $\|\gamma\| = \|\beta\|$ ($\|\gamma\| = \|\alpha\|$) for any $\gamma \in \Delta_s$ (respectively, for any $\gamma \in \Delta_l$), and by (13.22),

$$\sqrt{2x_2}|\beta|_1 = \|\alpha + 2\beta\| \leq \|\alpha + \beta\| + \|\beta\| = 2\|\beta\| = 2\sqrt{x_1}|\beta|_1,$$

which is equivalent to $x_2 \leq 2x_1$. Thus we get inequalities (13.21). ■

Remark 14. As it follows from the proof of inequalities (13.21), for a G - δ -homogeneous Riemannian manifold $(G/H, \mu)$ with $x_2 = 2x_1$ the restriction of the Chebyshev's norm $\|\cdot\|$ to the Cartan subalgebra $\mathfrak{t}$ is not strictly convex norm.

Corollary 14. *Every compact G - δ -homogeneous Riemannian flag manifold $M = G/T$ with a simple compact connected Lie group G is G -normal.*

Proof. Let suppose that the space under consideration is not G -normal. Then the first two statements in Proposition 33 imply that $C \neq \emptyset$. But this is impossible for $\mathfrak{h} = \mathfrak{t}$. ■

Proposition 34. *Every vector in $\mathfrak{p}_2$ for every G - δ -homogeneous Riemannian space $(G/H, \mu)$ is a δ -vector.*

Proof. Let's take in the above notation $t := x_1 \leq x_2$ and an arbitrary (non-zero) vector $v \in \mathfrak{p}_2$. Let suppose at first that $t = x_2$. In this case the corresponding space $M_2 = (G/H, \mu_2)$ is G -normal. Then there is unique Killing vector field on M_2 , which as an element of Lie algebra of right-invariant vector fields on G can be naturally identified with $v \in \mathfrak{p}_2 \subset \mathfrak{g}$. Then the Chebyshev's norm $\|X\|_2 = \sqrt{\mu_2(X(y), X(y))}$, where $y = H \in G/H = M$. Now, if we take $t = x_1 < x_2$, leaving x_2 fixed, then for any point $z \in M$ we will have

$$\sqrt{\mu(X(z), X(z))} \leq \sqrt{\mu_2(X(z), X(z))} \leq \sqrt{\mu_2(X(y), X(y))} = \sqrt{\mu(X(y), X(y))}.$$

This means that $y = H$ is a point of maximal distortion of X for μ also, which finishes the proof. ■

The following proposition follows from $\text{Ad}(G)$ -invariance of the Chebyshev's norm.

Proposition 35. *The set of all δ -vectors in some vector subspace $V_1 \subset \mathfrak{p}_1$ is invariant under all $\text{Ad}(g)$, $g \in G$, which leave V_1 invariant.*

14. THE SPECIAL SECOND CASE

Now we suppose that we have the second possibility in the Proposition 32, hence Δ contain roots of different length by Proposition 32 and $G \neq G_2$ by Section 11. So we need to consider only the simple Lie groups F_4 , and $Sp(l), SO(2l+1)$, when $l \geq 1$.

If $l = 1$, then the center $C(Sp(1))$ is isomorphic to $\mathbb{Z}_2$ and $Sp(1)/C(Sp(1)) = SO(3)$. The unique nontrivial Riemannian homogeneous space of positive Euler characteristic in this case is the symmetric (irreducible) space $Sp(1)/T = SO(3)/T = S^2$ of rank 1, which is G -normal, hence G - δ -homogeneous.

Proposition 36. *In the notation above, the following statements hold:*

- 1) *If $G \neq Sp(l)$, $l \geq 3$, then $A \cup C = \Delta_l$, $B = \Delta_s$.*
- 2) *If $G = Sp(l)$, $l \geq 3$, then for every $\alpha \in A$ and $\gamma \in C$, $\langle \alpha, \gamma \rangle = 0$ and $[V_\alpha, V_\gamma] = 0$.*
- 3) *For every $\alpha \in A$ there is an $\beta \in B$ such that $\langle \alpha, \beta \rangle \neq 0$. If $G \neq G_2$, then one (and only one) of the vectors $\alpha + \beta$ or $\alpha - \beta$ is root in B , and $\alpha + 2\beta$ (respectively, $\alpha - 2\beta$) is a root in C .*

Proof. The first statement in the case $G \neq F_4$ follows from the statement 2) of Lemma 7 and from the inclusion $\eta \subset \mathfrak{h} \oplus \mathfrak{p}_2$.

Suppose that $G = F_4$. By Lemma 8, $\mathfrak{p}_2 \oplus \mathfrak{h}$ is a proper Lie subalgebra in f_4 , which contains η by Lemma 7. So, by the statement 4) in Lemma 7, either $\mathfrak{p}_2 \oplus \mathfrak{h} = \eta$, or $\mathfrak{p}_2 \oplus \mathfrak{h} = \mathfrak{r}_i$ for some $1 \leq i \leq 3$. The second case is impossible. Suppose the contrary. Since $\mathfrak{r}_i = \eta \oplus \mathfrak{q}_i$, we get $\mathfrak{q}_i \subset \mathfrak{h}$. On the other hand, the module $\mathfrak{q}_i$ generates the Lie algebra $\mathfrak{r}_i$ ($(\mathfrak{r}_i, \eta) = (so(9), so(8))$). Since $\mathfrak{h}$ is a proper subalgebra in $\mathfrak{r}_i$, this is impossible. Therefore, $\mathfrak{p}_2 \oplus \mathfrak{h} = \eta$ and B coincides with the set Δ_s . This proves the first statement for $G = F_4$.

The second statement follows from the statement 3) of Lemma 7, if $\gamma \in \Delta_l$. The case $\gamma \in \Delta_s$, can be considered as Lemma 8 above.

Consider now the item 3). For any $\alpha \in A$ there is $\beta \in \Delta_s$ such that $\gamma := \alpha + \beta \in \Delta$ (otherwise an angle between α and any $\beta \in \Delta_s$ is $\pi/2$, with using the Weyl group we get the same for any root in A , but the latter contradicts to Lemma 5). It is clear that $\gamma \in \Delta_s$. Since $\gamma - \beta = \alpha \in A$, then either β or γ is not in C , hence one of them is in B . Other statements of this item follow from Lemma 8 and from Proposition 33. ■

Proposition 37. *Up to change of indices, in the case of $G = Sp(l)$, we must have $A = \{\pm 2e_1\}$, $\{\pm e_1 \pm e_i, 1 < i \leq l\} \subset B$.*

Proof. Let suppose that A contains besides $\pm 2e_1$ (up to change of indices) yet $\pm 2e_2$. Then by the statement 2) in Proposition 36, C cannot contain roots of the form $\pm e_i \pm e_j$, $i < j$, where $i = 1$ or $i = 2$. So, B contains all roots of the form $\pm e_1 \pm e_i$, $1 < i$, and $\pm e_2 \pm e_j$, $2 < j$. Let consider the root $-e_1 + e_2 \in B$. Then $[V_{2e_1}, V_{-e_1+e_2}] = V_{e_1+e_2}$. Now by Lemma 4

$$[V_{e_1+e_2}, V_{-e_1+e_2}] = V_{2e_2} \oplus V_{2e_1} \subset \mathfrak{p}_2.$$

So, in the previous notation

$$\alpha := 2e_1, \quad \beta := -e_1 + e_2, \quad \alpha + \beta = e_1 + e_2, \quad \alpha + 2\beta = 2e_2 \in A.$$

We have got a contradiction with the second part of the second statement in 3) of Proposition 36.

Now $A = \{\pm 2e_1\}$ and by the first part of the second statement in 3) of Proposition 36, all roots of the form $\pm e_1 \pm e_i$, $1 < i$, must lie in B . ■

Corollary 15. *In conditions of Proposition 37, $\dim(\mathfrak{p}_2) = 2$.*

Proposition 38. *For the case $G = Sp(l)$, $l \geq 2$, the spaces under consideration may have only one of the form $M = Sp(l)/U(1) \cdot Sp(l-1)$ or $Sp(l)/U(1) \times Sp(k_2-1) \times \cdots \times Sp(l-k_m)$, where $1 < k_2 < \cdots < k_m < l$, $m \geq 2$.*

Proof. In the Notation of Proposition 37, let suppose also that all other short roots (of the form $\pm e_i \pm e_j$, $2 \leq i < j \leq l$) lie in C . In this case we get exactly the first case. Here $U(1) \cdot Sp(l-1)$ is the centralizer of the root $2e_1 \in \mathfrak{t}$ and $\mathfrak{h} \oplus \mathfrak{p}_2 = sp(1) \oplus sp(l-1) \subset sp(l)$.

Let suppose that in the previous conditions $G = Sp(l)$ and $H \neq U(1) \times Sp(l-1)$. From Propositions 37 and the first case we get that

$$U(1) \times Sp(1)^{l-1} \subset H \subset U(1) \times Sp(l-1) \subset Sp(1) \times Sp(l-1).$$

Therefore, we obtain the second case from the description of subgroups with maximal rank of the group $Sp(l)$, obtained in Theorem II of [47]. ■

Theorem 27. *For the case $G = Sp(l)$, $l \geq 2$, the spaces under consideration may have only the form $M = Sp(l)/U(1) \cdot Sp(l-1)$.*

Proof. Suppose the contrary, then according to Proposition 38 there is a δ -homogeneous Riemannian manifold $(G/H = Sp(l)/U(1) \times Sp(k_2-1) \times \cdots \times Sp(l-k_m), \mu = \mu_{x_1, x_2})$, where $1 < k_2 < \cdots < k_m < l$, $m \geq 2$, and $x_1 \neq x_2$.

Let $K = Sp(1) \times Sp(k_2-1) \times \cdots \times Sp(l-k_m)$, $H \subset K \subset G$. Then $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1$, $sp(1) = u(1) \oplus \mathfrak{p}_2$. We will use notation $\mathfrak{h}_1 = u(1)$, $\mathfrak{h}_2 = sp(k_2-1) \oplus \cdots \oplus sp(l-k_m)$, where $\mathfrak{h} = \mathfrak{h}_1 \oplus \mathfrak{h}_2$. Let us consider $\text{Ad}(H)$ -invariant submodules $\mathfrak{p}_{1,1}, \mathfrak{p}_{1,2} \subset \mathfrak{p}_1$ such that $\mathfrak{g} = sp(l) = sp(1) \oplus sp(l-1) \oplus \mathfrak{p}_{1,1}$, $sp(l-1) = \mathfrak{h}_2 \oplus \mathfrak{p}_{1,2}$, where all sums are orthogonal with respect to $\langle \cdot, \cdot \rangle$, and $\mathfrak{p}_1 = \mathfrak{p}_{1,1} \oplus \mathfrak{p}_{1,2}$.

Take any $X \in \mathfrak{p}_{1,1} \subset \mathfrak{p}_1$ and any nontrivial $Y \in \mathfrak{p}_2$. Then there is some $Z \in \mathfrak{h}$ such that the vector $X + Y + Z$ is a δ -vector. In particular, this vector is geodesic for $(G/H, \mu)$. Then using Proposition 21 we get that $[Z, Y] = 0$. This means that $Z \in \mathfrak{h}_2$.

Take now any $U \in \mathfrak{p}_{1,2} \subset \mathfrak{p}_1$ and apply the inequality (10.10) from Proposition 22 in this situation. It is clear that $[U, X] \in \mathfrak{p}_{1,1} \subset \mathfrak{p}_1$, $[U, Y] = 0$, $[Z, U] \subset \mathfrak{p}_{1,2}$ and $\langle [U, X], [U, Z] \rangle = 0$. Hence the inequality (10.10) take the form $x_1 \langle [U, Z], [U, Z] \rangle \leq 0$, consequently, $[U, Z] = 0$ for any $U \in \mathfrak{p}_{1,2}$. On the other hand, it is easy to see, that the submodule $\mathfrak{p}_{1,2}$ generates the Lie algebra $sp(l-1)$ (the pair $(sp(l-1), \mathfrak{h}_2)$ is effective), therefore Z sits in the center of $sp(l-1)$ and $Z = 0$.

Now we use Proposition 21 again. We get that $[X, Y] = x_1/(x_2 - x_1)[X, Z] = 0$. Since $X \in \mathfrak{p}_{1,1}$ is arbitrary we get $[Y, \mathfrak{p}_{1,1}] = 0$. This is impossible since Y is nontrivial and the submodule $\mathfrak{p}_{1,1}$ generates the Lie algebra $sp(l)$. Therefore, $(G/H, \mu)$ is not δ -homogeneous. Theorem is proved. ■

Theorem 28. *If $G = SO(2l + 1)$, where $l \geq 2$, then the space $M := G/H$ under consideration may have only one form $M = SO(2l + 1)/U(l)$.*

Proof. The group $G = SO(2l + 1)$ has the root system B_l . Then the Lie algebra η from Lemma 7 is isomorphic to the Lie algebra $so(2l)$ of the Lie group $SO(2l)$ with the root system D_l . In this case $\eta = \mathfrak{h} \oplus \mathfrak{p}_2$ and $\mathfrak{p}_1 = \text{Lin}\{\cup_{\beta \in \Delta_s} V_\beta\}$ by the statement 1) in Proposition 36. Therefore the homogeneous space $(SO(2l + 1)/H, \mu)$ under consideration is fibred over rank 1 (hence irreducible) symmetric space $SO(2l + 1)/SO(2l) = S^{2l}$. So, the conditions of Theorem 4.1 in the paper [42] are satisfied. Then by Table I on the page 841 of this paper and by Theorem 17 we must have $M = SO(2l + 1)/U(l)$. ■

Remark 15. The spaces in Theorem 27 and the spaces from Theorem 28 were appeared also in the paper [4] as (generalized) flag manifolds, admitting non-normal invariant g.o. Riemannian metrics. Earlier W. Ziller proved that all invariant metrics on these spaces are weakly symmetric (hence, g.o.) [52].

Corollary 16. *For spaces in Theorem 28, every vector in $\mathfrak{p}_1$ is a δ -vector.*

Proof. By the proof of Theorem 28, $\mathfrak{p}_1$ is naturally identified with the tangent space at the initial point of a rank 1 symmetric space $SO(2l + 1)/SO(2l) = S^{2l}$, which is two-point homogeneous. This implies that $\text{Ad}(SO(2l + 1))$ acts transitively on the unit sphere in $(\mathfrak{p}_1, (\cdot, \cdot))$. The proof is finished by applying of Propositions 31 and 35. ■

Theorem 29. *There is no space $M := G/H$ under consideration with $G = F_4$.*

Proof. At first, we prove that $M = G/H$ with $G = F_4$ may have at most one form $M = F_4/\exp(u(4))$.

In this case $\mathfrak{h} \oplus \mathfrak{p}_2 = \eta = so(8)$, $\mathfrak{p}_1 = \mathfrak{q}_1 \oplus \mathfrak{q}_2 \oplus \mathfrak{q}_3$ (see Lemma 7 and Proposition 36). Let's consider a subalgebra $\mathfrak{r}_3 = \eta \oplus \mathfrak{q}_3 = so(9) = spin(9) = \mathfrak{l}$. By Proposition 9, the Riemannian subspace $L/H = Spin(9)/H \subset F_4/H$ is totally geodesic, hence δ -homogeneous and g.o. space, and also has positive Euler characteristic. Since $L = Spin(9)$ is a simple group and the restriction of the Killing form of f_4 to $\mathfrak{l}$ is $\text{Ad}(L)$ -invariant, then this restriction must be proportional to the Killing form of $\mathfrak{l}$. We have $Spin(9)/H = (Spin(9)/C)/(H/C) = SO(9)/(H/C)$, where C is the common center of $Spin(9)$ and H . Therefore, the Riemannian subspace $SO(9)/(H/C)$ of F_4/H is not $SO(9)$ -normal, if F_4/H is not F_4 -normal, because $\mathfrak{l}$ includes vector subspaces $\mathfrak{p}_2$ and $\mathfrak{q}_3 \subset \mathfrak{p}_1$.

If $SO(9)/(H/C)$ is not $SO(9)$ - δ -homogeneous (being δ -homogeneous), then its full connected isometry group is not equal to $SO(9)$. Therefore, according to Theorem 24, we must have $H/C = U(4)$ and $H = \exp(u(4))$. On the other hand, if $SO(9)/(H/C)$ is $SO(9)$ - δ -homogeneous, then by Theorem 28, we get again $H = \exp(u(4))$. Note that $\mathfrak{h} = u(4)$ is spanned on the Cartan subalgebra $\mathfrak{t}$ and on the root spaces of the roots $\pm(e_i - e_j)$, $1 \leq i < j \leq 4$.

Now we shall prove that the Riemannian manifold $(G/H = F_4/\exp(u(4)), \mu = \mu_{x_1, x_2})$ is not g.o. for $x_1 \neq x_2$.

Note that the submodule $\mathfrak{q}_2$ (see Lemma 7) is not $\text{ad}(\mathfrak{h})$ -irreducible. Really, let us consider a two-dimensional submodule $\mathfrak{q} \subset \mathfrak{q}_2$, which is spanned on the root space of the vectors $\pm 1/2(e_1 + e_2 + e_3 + e_4)$. It is clear that

$$(\pm(e_i - e_j)) + (\pm 1/2(e_1 + e_2 + e_3 + e_4))$$

is not a root for any $1 \leq i < j \leq 4$. This means that $\mathfrak{q}$ commutes with every root spaces of the roots $\pm(e_i - e_j)$. Therefore, $\mathfrak{q}$ is invariant under the action of $\text{ad}(\mathfrak{h})$.

Consider now any $X \in \mathfrak{q} \subset \mathfrak{q}_2 \subset \mathfrak{p}_1$ and any $Y \in \mathfrak{p}_2$. If $(F_4/\exp(u(4)), \mu)$ is a g.o. space, then there is $Z \in \mathfrak{h}$ such that $X + Y + Z$ is a geodesic vector. If we have $x_1 \neq x_2$ in addition, then according to Proposition 21, we get $[X, Y] = x_1/(x_2 - x_1)[X, Z]$. Since $[X, Z] \subset \mathfrak{q}$, we obtain that $[X, Y] \in \mathfrak{q}$ for any $X \in \mathfrak{q}$ and for any $Y \in \mathfrak{p}_2$. Therefore, the module $\mathfrak{q}$ is $\text{ad}(\eta)$ -invariant which is impossible, since the module $\mathfrak{q}_2$ (containing $\mathfrak{q}$) is $\text{ad}(\eta)$ -irreducible. Therefore, $(F_4/\exp(u(4)), \mu)$ is not g.o. for $x_1 \neq x_2$. This finishes the proof. ■

15. ON THE SPACE $SO(5)/U(2) = Sp(2)/U(1) \cdot Sp(1) = \mathbb{C}P^3$

Here we find all δ -homogeneous metrics on the space $SO(5)/U(2)$, where $U(2) \subset SO(4) \subset SO(5)$, and the pair $(SO(5), SO(4))$, $(SO(4), U(2))$ are irreducible symmetric. Remind that the space $SO(5)/U(2)$ coincides with the space $Sp(2)/U(1) \cdot Sp(1)$.

For $A, B \in so(5)$ we define $\langle A, B \rangle = -1/2 \text{trace}(A \cdot B)$. This is an $\text{Ad}(SO(5))$ -invariant inner product on $so(5)$. A matrix $A + \sqrt{-1}B \in u(2)$, where $A = \begin{pmatrix} 0 & c \\ -c & 0 \end{pmatrix}$ and $B = \begin{pmatrix} a & d \\ d & b \end{pmatrix}$ we embed into $so(4)$ via $A + \sqrt{-1}B \mapsto \begin{pmatrix} A & B \\ -B & A \end{pmatrix}$ in order to get the symmetric pair $(so(4), u(2))$ (see e.g. [23]). Also we use the standard embedding $so(4)$ into $so(5)$: $A \mapsto \text{diag}(A, 0)$.

It is known the following $\langle \cdot, \cdot \rangle$ -orthogonal decomposition:

$$\mathfrak{g} = so(5) = so(4) \oplus \mathfrak{p}_1 = u(2) \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_1, \quad \mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2,$$

where

$$u(2) = \left\{ \begin{pmatrix} 0 & c & a & d & 0 \\ -c & 0 & d & b & 0 \\ -a & -d & 0 & c & 0 \\ -d & -b & -c & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} ; \quad a, b, c, d \in \mathbb{R} \right\},$$

$$\mathfrak{p}_1 = \left\{ X = \begin{pmatrix} 0 & 0 & 0 & 0 & k \\ 0 & 0 & 0 & 0 & l \\ 0 & 0 & 0 & 0 & m \\ 0 & 0 & 0 & 0 & n \\ -k & -l & -m & -n & 0 \end{pmatrix} ; \quad k, l, m, n \in \mathbb{R} \right\},$$

$$\mathfrak{p}_2 = \left\{ Y = \begin{pmatrix} 0 & e & 0 & f & 0 \\ -e & 0 & -f & 0 & 0 \\ 0 & f & 0 & -e & 0 \\ -f & 0 & e & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} ; \quad e, f \in \mathbb{R} \right\},$$

and the modules $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are $\text{Ad}(U(2))$ -invariant and $\text{Ad}(U(2))$ -irreducible. Note that for vectors X from $\mathfrak{p}_1$ as above we have $\langle X, X \rangle = k^2 + l^2 + m^2 + n^2$, and for vectors $Y \in \mathfrak{p}_2$ we have $\langle Y, Y \rangle = 2e^2 + 2f^2$.

Let us consider the invariant metric $\mu = \mu_{x_1, x_2}$ on $SO(5)/U(2)$, corresponding to the inner product

$$\langle \cdot, \cdot \rangle = x_1 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_1} + x_2 \langle \cdot, \cdot \rangle|_{\mathfrak{p}_2}$$

for some positive x_1 and x_2 . We know that every such metric is a g.o.-metric [52],[42]. From the discussion in Section 8 we get the following

Proposition 39. *The full connected isometry group of $(SO(5)/U(2), \mu)$ is $SO(5)$, excepting the case $x_2 = 2x_1$, where the full connected isometry group is $SO(6)/\{\pm I\}$, and the metric μ is $SO(6)$ -normal (in the last case $(SO(5)/U(2), \mu)$ is isometric to the complex projective space $\mathbb{C}P^3$ with the standard Fubini metric).*

Let $E_{i,j}$ be a (5×5) -matrix, whose (i, j) -th entry is equal to 1, and all other entries are zero. For any $1 \leq i < j \leq 5$ put $F_{i,j} = E_{i,j} - E_{j,i}$. Let consider the following subspace of $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$:

$$\mathfrak{q} = \mathbb{R} \cdot F_{1,5} \oplus \mathbb{R} \cdot (F_{1,4} - F_{2,3}).$$

Proposition 40. *For any vector $V \in \mathfrak{p}$ there is $a \in H = U(2)$ such that $\text{Ad}(a)(V) \in \mathfrak{q}$.*

Proof. Let $V = X + Y$, where $X \in \mathfrak{p}_1$ and $Y \in \mathfrak{p}_2$. We know by (the proof of) Corollary 16 that $\text{Ad}(U(2))$ acts transitively on the unit sphere in $\mathfrak{p}_1$. Therefore, we may assume that $X = bF_{1,5}$ for some $b \in \mathbb{R}$. We have

$$Y = c_1(F_{1,2} - F_{3,4}) + c_2(F_{1,4} - F_{2,3})$$

for some real c_1 and c_2 . Note that $[F_{2,4}, X] = 0$. Therefore, X is invariant under $\text{Ad}(a)$, where $a = \exp(tF_{2,4})$. On the other hand,

$$\text{Ad}(a)(Y) = \tilde{c}_1(F_{1,2} - F_{3,4}) + \tilde{c}_2(F_{1,4} - F_{2,3}) \in \mathfrak{p}_2,$$

where

$$\tilde{c}_1 = c_1 \cos(t) + c_2 \sin(t), \quad \tilde{c}_2 = c_2 \cos(t) - c_1 \sin(t).$$

For some suitable $t \in \mathbb{R}$ we get that $\tilde{c}_1 = 0$. Therefore, $\text{Ad}(a)(V) = bF_{1,5} + \tilde{c}_2(F_{1,4} - F_{2,3}) \in \mathfrak{q}$. $\blacksquare$

Proposition 41. *Let $W = X + Y + Z$, where $X + Y \in \mathfrak{q}$ and $Z \in \mathfrak{h} = u(2)$, be a non-trivial geodesic vector on $(SO(5)/U(2), \mu)$, $x_2 \neq x_1$, $x_2 \neq 2x_1$. Then we have one of the following possibilities:*

- 1) $W = bF_{1,5} + \frac{x_2}{x_1}cF_{1,4} + \frac{x_2-2x_1}{x_1}cF_{2,3}$ for some $b \neq 0$, $c \neq 0$;
- 2) $W = d(F_{1,4} - F_{2,3}) + a_1(F_{1,2} + F_{3,4}) + a_2(F_{1,4} + F_{2,3}) + a_3(F_{1,3} - F_{2,4})$ for some $d \neq 0$, $a_1, a_2, a_3 \in \mathbb{R}$;
- 3) $W = eF_{1,5} + fF_{2,4}$ for some $e \neq 0$ and $f \in \mathbb{R}$.

Proof. Let $W = X + Y + Z$, where $X = bF_{1,5} \in \mathfrak{p}_1$, $Y = c(F_{1,4} - F_{2,3})$, and $Z = b_1(F_{1,2} + F_{3,4}) + b_2(F_{1,4} + F_{2,3}) + b_3F_{1,3} + b_4F_{2,4}$. Since W is geodesic vector, then from Proposition 21 we have

$$[Z, Y] = 0, \quad [X, Y] = \frac{x_1}{x_2 - x_1}[X, Z].$$

Direct calculations show that

$$[Z, Y] = c(b_3 + b_4)(F_{1,2} - F_{3,4}), \quad [X, Y] = bcF_{4,5}, \quad [X, Z] = b(b_1F_{2,5} + b_3F_{3,5} + b_2F_{4,5}).$$

If $b \neq 0$ and $c \neq 0$, then $b_1 = b_3 = b_4 = 0$ and $b_2 = \frac{x_2 - x_1}{x_1}c$.

If $b = 0$ and $c \neq 0$, then $b_4 = -b_3$.

If $b \neq 0$ and $c = 0$, then we have $b_1 = b_2 = b_3 = 0$.

The proposition is proved. $\blacksquare$

Proposition 42. *The Riemannian manifold $(SO(5)/U(2), \mu)$ is $SO(5) - \delta$ -homogeneous if and only if for every $b \neq 0$ and every $c \neq 0$ the vector*

$$W = \begin{pmatrix} 0 & 0 & 0 & \frac{x_2}{x_1}c & b \\ 0 & 0 & \frac{x_2-2x_1}{x_1}c & 0 & 0 \\ 0 & \frac{2x_1-x_2}{x_1}c & 0 & 0 & 0 \\ -\frac{x_2}{x_1}c & 0 & 0 & 0 & 0 \\ -b & 0 & 0 & 0 & 0 \end{pmatrix} = bF_{1,5} + \frac{x_2}{x_1}cF_{1,4} + \frac{x_2-2x_1}{x_1}cF_{2,3}$$

is δ -vector on $(SO(5)/U(2), \mu)$.

Proof. If $(SO(5)/U(2), \mu)$ is $SO(5) - \delta$ -homogeneous, then for every vector of the form $V = X + Y$, where $X = bF_{1,5} \in \mathfrak{p}_1$, $Y = c(F_{1,4} - F_{2,3}) \in \mathfrak{p}_2$, $b \neq 0$, $c \neq 0$, there is $Z \in \mathfrak{h}$ such that the vector $W = X + Y + Z$ is δ -vector. In particular, W is geodesic vector. According to Proposition 41, we get that

$$W = bF_{1,5} + \frac{x_2}{x_1}cF_{1,4} + \frac{x_2-2x_1}{x_1}cF_{2,3}.$$

Therefore, this W is a δ -vector.

Let us suppose now that all vectors of the form

$$W = bF_{1,5} + \frac{x_2}{x_1}cF_{1,4} + \frac{x_2-2x_1}{x_1}cF_{2,3},$$

where $b \neq 0$ and $c \neq 0$, are δ -vectors. Since the limit of any sequence of δ -vectors is a δ -vector itself, we get that the vectors W as above are δ -vectors for $b = 0$ or $c = 0$ also.

Therefore, for any vector $X + Y \in \mathfrak{q}$ there is $Z \in \mathfrak{h}$ such that the vector $X + Y + Z$ is δ -vector. Using Proposition 40, we get that $(SO(5)/U(2), \mu)$ is $SO(5)$ - δ -homogeneous in this case. ■

Lemma 9. *For every $b, c, x_1, x_2 \in \mathbb{R}$ with the properties*

$$b \neq 0, \quad x_1 \neq 0, \quad 2x_1 > x_2,$$

the following inequality is fulfilled:

$$\left(|c|(2x_1 - x_2) + \sqrt{b^2x_1^2 + c^2x_2^2} \right)^2 x_2 < 2x_1^2(x_1b^2 + 2x_2c^2).$$

Proof. It is enough to consider the case $x_2 > 0$. In this case we have the following chain of equivalent inequalities.

$$\begin{aligned} & \left(c^2(2x_1 - x_2)^2 + b^2x_1^2 + c^2x_2^2 + 2|c|(2x_1 - x_2)\sqrt{b^2x_1^2 + c^2x_2^2} \right) x_2 < 2x_1^3b^2 + 4x_1^2x_2c^2; \\ & 2|c|(2x_1 - x_2)\sqrt{b^2x_1^2 + c^2x_2^2}x_2 < 2x_1^3b^2 + 4x_1^2x_2c^2 - c^2(2x_1 - x_2)^2x_2 - b^2x_1^2x_2 - c^2x_2^3 = \\ & \quad (2x_1 - x_2)x_1^2b^2 + 2x_2^2(2x_1 - x_2)c^2; \\ & 2|c|\sqrt{b^2x_1^2 + c^2x_2^2}x_2 < x_1^2b^2 + 2x_2^2c^2; \\ & 4c^2(b^2x_1^2 + c^2x_2^2)x_2^2 = 4x_1^2x_2^2b^2c^2 + 4x_2^4c^4 < x_1^4b^4 + 4x_1^2x_2^2b^2c^2 + 4x_2^4c^4 = (x_1^2b^2 + 2x_2^2c^2)^2. \end{aligned}$$

■

Proposition 43. *If $2x_1 \geq x_2 \geq x_1$, then the Riemannian manifold $(SO(5)/U(2), \mu)$ is $SO(5)$ - δ -homogeneous.*

Proof. We may assume by continuity, that $x_1 < x_2 < 2x_1$.

According to Proposition 42, we only need to prove that every vector of the form

$$W = bF_{1,5} + \frac{x_2}{x_1}cF_{1,4} + \frac{x_2 - 2x_1}{x_1}cF_{2,3},$$

where $b \neq 0$ and $c \neq 0$, is δ -vector on $(SO(5)/U(2), \mu)$.

Let us consider the orbit $O(W)$ of W under the action of $\text{Ad}(G) = \text{Ad}(SO(5))$. Since $O(W)$ is compact, there is $\widetilde{W} \in O(W)$ such that

$$(\widetilde{W}|_{\mathfrak{p}}, \widetilde{W}|_{\mathfrak{p}}) \geq (V|_{\mathfrak{p}}, V|_{\mathfrak{p}})$$

for every $V \in O(W)$.

Therefore, $\widetilde{W}$ is a δ -vector. According to Proposition 40 we may assume, that $\widetilde{W}|_{\mathfrak{p}} \in \mathfrak{q}$. Now it is sufficient to show that

$$(\widetilde{W}|_{\mathfrak{p}}, \widetilde{W}|_{\mathfrak{p}}) \leq (W|_{\mathfrak{p}}, W|_{\mathfrak{p}})$$

We shall use the following idea. Since $\widetilde{W} \in O(W)$, then the matrices $-W^2$ and $-\widetilde{W}^2$ has one and the same set of eigenvalues. The eigenvalues of $-W^2$ are the following:

$$0, \quad \frac{c^2(2x_1 - x_2)^2}{x_1^2}, \quad \frac{b^2x_1^2 + c^2x_2^2}{x_1^2},$$

where two last eigenvalues are of multiplicity 2. Since $x_2 > x_1$, we obviously get

$$b^2x_1^2 + c^2x_2^2 > c^2(2x_1 - x_2)^2.$$

Note also that $(W|_{\mathfrak{p}}, W|_{\mathfrak{p}}) = x_1b^2 + 2x_2c^2$.

Since $\widetilde{W}$ is geodesic vector and $\widetilde{W}|_{\mathfrak{p}} \in \mathfrak{q}$, then by Proposition 41 we have one of the following possibilities:

$$1) \quad \widetilde{W} = \widetilde{b}F_{1,5} + \frac{x_2}{x_1}\widetilde{c}F_{1,4} + \frac{x_2 - 2x_1}{x_1}\widetilde{c}F_{2,3} \text{ for some } \widetilde{b} \neq 0, \widetilde{c} \neq 0;$$

2) $\widetilde{W} = d(F_{1,4} - F_{2,3}) + a_1(F_{1,2} + F_{3,4}) + a_2(F_{1,4} + F_{2,3}) + a_3(F_{1,3} - F_{2,4})$ for some $d \neq 0$, $a_1, a_2, a_3 \in \mathbb{R}$;

3) $\widetilde{W} = eF_{1,5} + fF_{2,4}$ for some $e \neq 0$ and $f \in \mathbb{R}$.

Let us consider these cases separately.

Case 1). Eigenvalues of $-\widetilde{W}^2$ in this case are the following:

$$0, \quad \frac{\tilde{c}^2(2x_1 - x_2)^2}{x_1^2}, \quad \frac{\tilde{b}^2x_1^2 + \tilde{c}^2x_2^2}{x_1^2},$$

where two last eigenvalues are of multiplicity 2. Since $\tilde{b}^2x_1^2 + \tilde{c}^2x_2^2 > \tilde{c}^2(2x_1 - x_2)^2$ (remind that $x_2 > x_1$) and $\widetilde{W} \in O(W)$, we get that

$$\tilde{b}^2x_1^2 + \tilde{c}^2x_2^2 = b^2x_1^2 + c^2x_2^2, \quad \tilde{c}^2(2x_1 - x_2)^2 = c^2(2x_1 - x_2)^2,$$

which implies $c^2 = \tilde{c}^2$ and $b^2 = \tilde{b}^2$ (since $2x_1 > x_2$). Therefore

$$(\widetilde{W}|_{\mathfrak{p}}, \widetilde{W}|_{\mathfrak{p}}) = x_1\tilde{b}^2 + 2x_2\tilde{c}^2 = x_1b^2 + 2x_2c^2 = (W|_{\mathfrak{p}}, W|_{\mathfrak{p}}).$$

Case 2). In this case the eigenvalues of $-\widetilde{W}^2$ are the following:

$$0, \quad d^2 + a_1^2 + a_2^2 + a_3^2 - 2\sqrt{d^2(a_1^2 + a_2^2 + a_3^2)}, \quad d^2 + a_1^2 + a_2^2 + a_3^2 + 2\sqrt{d^2(a_1^2 + a_2^2 + a_3^2)},$$

where two last eigenvalues are of multiplicity 2.

Since $\widetilde{W} \in O(W)$, we obtain

$$(|d| - |s|)^2 = d^2 + s^2 - 2\sqrt{d^2s^2} = \frac{c^2(2x_1 - x_2)^2}{x_1^2}, \quad (|d| + |s|)^2 = d^2 + s^2 + 2\sqrt{d^2s^2} = \frac{b^2x_1^2 + c^2x_2^2}{x_1^2},$$

where $s^2 = a_1^2 + a_2^2 + a_3^2$. We get from these equations

$$2|d| = (|d| - |s|) + (|d| + |s|) \leq \frac{|c|(2x_1 - x_2)}{x_1} + \frac{\sqrt{b^2x_1^2 + c^2x_2^2}}{x_1}.$$

Using Lemma 9, we get

$$4d^2x_1x_2 \leq \left(|c|(2x_1 - x_2) + \sqrt{b^2x_1^2 + c^2x_2^2} \right)^2 x_2 < 2x_1^2(x_1b^2 + 2x_2c^2).$$

Therefore

$$(\widetilde{W}|_{\mathfrak{p}}, \widetilde{W}|_{\mathfrak{p}}) = 2x_2d^2 < x_1b^2 + 2x_2c^2 = (W|_{\mathfrak{p}}, W|_{\mathfrak{p}}).$$

Case 3). In this case the eigenvalues of $-\widetilde{W}^2$ are the following:

$$0, \quad e^2, e^2, f^2, f^2.$$

Therefore

$$e^2 = \frac{c^2(2x_1 - x_2)^2}{x_1^2} \quad \text{or} \quad e^2 = \frac{b^2x_1^2 + c^2x_2^2}{x_1^2},$$

Since $2x_1 > x_2 > x_1$, we get

$$x_1b^2 + 2x_2c^2 > \frac{b^2x_1^2 + c^2x_2^2}{x_1} > \frac{c^2(2x_1 - x_2)^2}{x_1},$$

which implies $x_1b^2 + 2x_2c^2 > x_1e^2$. Therefore

$$(\widetilde{W}|_{\mathfrak{p}}, \widetilde{W}|_{\mathfrak{p}}) = x_1e^2 < x_1b^2 + 2x_2c^2 = (W|_{\mathfrak{p}}, W|_{\mathfrak{p}}).$$

The above considerations prove that W is a δ -vector on $(SO(5)/U(2), \mu)$. This proves the proposition. ■

Theorem 30. *The Riemannian manifold $(SO(5)/U(2), \mu = \mu_{x_1, x_2})$ is δ -homogeneous if and only if $x_1 \leq x_2 \leq 2x_1$. For $x_2 = x_1$ it is $SO(5)$ -normal homogeneous; for $x_2 = 2x_1$ it is $SO(6)$ -normal homogeneous; for $x_2 \in (x_1, 2x_1)$ it is not normal homogeneous with respect to any its isometry group, but $SO(5)$ - δ -homogeneous.*

Proof. If $(SO(5)/U(2), \mu = \mu_{x_1, x_2})$ is δ -homogeneous, then it is $SO(6)$ - δ -homogeneous or $SO(5)$ - δ -homogeneous, see Theorem 23. In the first case it is $SO(6)$ -homogeneous. Then by Example 6, we have $x_2 = 2x_1$. In the second case, by Proposition 33 we get $x_1 \leq x_2 \leq 2x_1$. On the other hand, for $x_2 = x_1$ and for $x_2 = 2x_1$ the metric μ is $SO(5)$ -normal homogeneous and $SO(6)$ -normal homogeneous respectively (see Example 6). From Proposition 43 we get that the Riemannian manifold $(SO(5)/U(2), \mu)$ is δ -homogeneous for $2x_1 > x_2 > x_1$. The theorem is proved. ■

Remark 16. According to Theorem 25, the metrics in Theorem 30 with the condition $x_2 \in (x_1, 2x_2)$ are not naturally reductive (with respect to any isometry group) in spite of the fact that they are δ -homogeneous.

Remark 17. It follows from [45] that the Riemannian manifolds in Theorem 30 have positive sectional curvatures and their (exact) pinch constant is $\varepsilon = (\frac{x_2}{4x_1})^2$. This means that if we scale them so that their maximal sectional curvature will be 1, then minimal sectional curvature will be ε .

REFERENCES

- [1] Adams, J. Frank. Lectures on Lie groups. W.A.Benjamin, Inc., New York-Amsterdam, 1969.
- [2] Adams, J. Frank. Lectures on exceptional Lie groups. The University of Chicago Press, Chicago-London, 1996.
- [3] Aleksevskii D.V. Compact quaternion spaces. Funk. Anal. Pril., **2(2)** (1968), 11–20.
- [4] Alekseevsky, D.V. and Arvanitoyeorgos, A. Riemannian flag manifolds with homogeneous geodesics, to appear in Trans. AMS.
- [5] Akhiezer D.N and Vinberg E.B. Weakly symmetric spaces and spherical varieties. Transformation Groups, **4** (1999), 3–24.
- [6] Berestovskii, V.N. Homogeneous Busemann G -spaces. (Russian) Sibirsk. Mat. Zh., **23 (2)** (1982), 3–15.
- [7] Berestovskii, V.N. Homogeneous Riemannian manifolds of positive Ricci curvature. (Russian) Mat. Zametki, **58 (3)** (1995), 334–340; translation in Math. Notes, **58 (3-4)** (1996), 905–909.
- [8] Berestovskii, Valera, and Plaut, Conrad. Homogeneous spaces of curvature bounded below. J. Geom. Anal., **9 (2)** (1999), 203–219.
- [9] Berestovskii, V.N. and Guijarro, Luis. A metric characterization of Riemannian submersions. Ann. Global Anal. Geom., **18 (6)** (2000), 577–588.
- [10] Berger, M. Les varietes riemanniennes homogenes normales a courbure strictement positive. Ann. Sc. Norm. Sup. Pisa, **15** (1961), 179–246.
- [11] Berger, M. Trois remarques sur les variétés riemanniennes à courbure positive. C. R. Acad. Sci. Paris Sér. A-B, **263** (1966), 76–78.
- [12] Besse, A.L. Einstein Manifolds. Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1987.
- [13] Borel, A. and de Siebenthal, J., Les sous-groupes fermés de rang maximum des groupes de Lie clos. Comment. Math. Helv., **23** (1949), 200–221.
- [14] Bourbaki, N. Groupes et algèbres de Lie. Chapter 9. Groupes de Lie réels compacts. MASSON, Paris, 1982.
- [15] Bott, R. Vector fields and characteristic numbers. Michigan Math. J., **14** (1967), 231–244.
- [16] Busemann, Herbert. The geometry of geodesics. Academic Press Inc., New York, N.Y., 1955.
- [17] Cheeger, J. and Gromoll, D. The splitting theorem for manifolds of nonnegative Ricci curvature. J. Dif. Geometry, **6(6)** (1971), 119–128.
- [18] Cohn-Vossen, S. Existenz kürzester Wege. Compositio Math., **3** (1936), 441–452.
- [19] D'Atri, J.E. and Ziller, W. Naturally reductive metrics and Einstein metrics on compact Lie groups. Memoirs Amer. Math. Soc., **18** (1979), no. 215, 1–72.
- [20] Dušek Z., Kowalski O., Nikčević S. Ž. New examples of Riemannian g.o. manifolds in dimension 7. Differential Geom. Appl., **21** (2004), 65–78.
- [21] Freudenthal, H. Clifford-Wolf-Isometrien symmetrischer Räume, Math. Ann., **150** (1963), 136–149.
- [22] Gorbatsevich, V.V. and Onishchik, A.L. Lie transformation groups. Lie groups and Lie algebras, I, 95–235. Encyclopedia Math. Sci., 20, Springer, Berlin, 1993.
- [23] Helgason, S. Differential geometry and symmetric spaces. Academic Press Inc., New-York, 1962.
- [24] Hopf, H. and Samelson, H. Ein Satz über die Wirkungräume geschlossener Liescher Gruppen. (German). Comment. Math. Helv., **13** (1940-41), 240–251.
- [25] Jacobson, Nathan. Lie algebras. Interscience Tracts in Pure and Applied Mathematics, No. 10. Interscience Publishers (a division of John Wiley and Sons), New York-London, 1962.

- [26] Kerr, M. Some new homogeneous Einstein metrics on symmetric spaces. *Trans. Amer. Math. Soc.*, **348** (1996), 153–171.
- [27] Kobayashi, Sh. Transformation groups in differential geometry. Springer, Berlin, 1972.
- [28] Kobayashi, S. and Nomizu, K. Foundations of differential geometry. Vol. I – A Wiley-Interscience Publication, New York, 1963; Vol. II – A Wiley-Interscience Publication, New York, 1969.
- [29] Kostant, B. On differential geometry and homogeneous spaces. *Proc. Math. Acad. Sci. USA*, **42** (1956), 258–261, 354–357.
- [30] Kostant, B. On holonomy and homogeneous spaces. *Nagoya Math. J.*, **12** (1957), 31–54.
- [31] Kowalski, O. and Vanhecke, L. Riemannian manifolds with homogeneous geodesics. *Boll. Unione Mat. Ital.* vii.Ser. B, **5** (1) (1991), 189–246.
- [32] Krämer M. Sphärische Untergruppen in kompakten zusammenhängenden Liegruppen. *Compositio Math.*, **38**(2) (1979), 129–153.
- [33] Lauret J. Commutative spaces which are not weakly symmetric. *Bull. London Math. Soc.*, **30** (1998), 29–36.
- [34] Mimura, M. and Toda, H. Topology of Lie groups, I and II. AMS Translations 91, Providence, Rhode Island, 1991.
- [35] O’Neill, B. The fundamental equations of a submersion, *Michigan Math. J.*, **13**(4) (1966), 459–469.
- [36] Nguyễn H. D. Compact weakly symmetric spaces and spherical pairs. *Proc. AMS*, **128**(11) 2000, 3425–3433.
- [37] Onishchik, A.L. Topology of Transitive Transformation Groups. Johann Ambrosius Barth: Leipzig, Berlin, Heidelberg, 1994.
- [38] Onishchik, A.L. The group of isometries of a compact Riemannian homogeneous space. *Differential geometry and its applications* (Eger, 1989), 597–616, *Colloq. Math. Soc. János Bolyai*, 56, North-Holland, Amsterdam, 1992.
- [39] Selberg A. Harmonic analysis and discontinuous groups in weakly symmetric spaces with applications to Dirichlet series. *J. Indian Math. Soc. N.S.*, **20** (1956), 47–87.
- [40] Shchetinin, A.N. On a class of compact homogeneous spaces. I. *Ann. Global Anal. Geom.*, **6** (1988), 119–140.
- [41] Shchetinin, A.N. On a class of compact homogeneous spaces. II. *Ann. Global Anal. Geom.*, **8** (1990), 227–247.
- [42] Tamaru, H. Riemannian g.o. spaces fibered over irreducible symmetric spaces. *Osaka J. Math.*, **36** (1999), 835–851.
- [43] Tamaru, H. Riemannian geodesic orbit metrics on fiber bundles. *Algebra, Groups and Geometries*, **15** (1998), 55–67.
- [44] Toponogov, V.A. Metric structure of Riemannian spaces of nonnegative curvature containing direct lines (Russian). *Sib. Math. Zh.*, **5**(5) (1964), 1358–1369.
- [45] Vol’per, D.E. Sectional curvatures of nonstandard metrics on CP^{2n+1} (Russian). *Sibirsk. Mat. Zh.*, **40**(1) (1999), 49–56; translation in *Siberian Math. J.*, **40**(1) (1999), 39–45.
- [46] Wallach, N.R. Compact homogeneous Riemannian manifolds with strictly positive curvature, *Annals of Math.*, **96**(2) (1972), 277–295.
- [47] Wang, H.C. Homogeneous spaces with non-vanishing Euler characteristic. *Annals of Math.*, 2nd Ser., **50**(4) (1949), 925–953.
- [48] Wang, M. and Ziller, W. On isotropy irreducible Riemannian manifolds. *Acta Math.*, **166** (1991), 223–261.
- [49] Warner, Frank W. Foundations of differentiable manifolds and Lie groups. Corrected reprint of the 1971 edition. Graduate Texts in Mathematics, 94. Springer-Verlag, New York-Berlin, 1983.
- [50] J.A. Wolf, Spaces of constant curvature. University of California, Berkley, 1972.
- [51] Ziller, W. Homogeneous Einstein Metrics on Spheres and Projective Spaces. *Math. Ann.*, **259** (1982), 351–358.
- [52] Ziller, W. Weakly symmetric spaces, 355–368. In: *Progress in Nonlinear Differential Equations. V. 20. Topics in geometry: in memory of Joseph D’Atri*. Birkhäuser, 1996.

BERESTOVSKIĬ VALERIĬ NIKOLAEVICH

OMSK BRANCH OF SOBOLEV INSTITUTE OF MATHEMATICS SD RAS
644099, OMSK, UL. PEVTSOVA, 13, RUSSIA

E-mail address: berestov@ofim.oscsbras.ru

NIKONOROV YURIĬ GENNADIEVICH

RUBTSOVSK INDUSTRIAL INSTITUTE
OF ALTAI STATE TECHNICAL UNIVERSITY AFTER I.I. POLZUNOV
658207, RUBTSOVSK, TRAKTORNAYA, 2/6, RUSSIA

E-mail address: nik@inst.rubtsovsk.ru